Lophotrochozoa

Members have either a **lophophore** (a crown of ciliated tentacles)

- **Bryozoans**
- **Brachiopods**
- **Rotifers**

Or have a **trochophore larval stage**

- Platyhelminthes (flatworms)
- Annelids
- **Molluscs**

PHYLUM ROTIFERA - “WHEEL ANIMALCULES”

- Named for ciliated crown or corona
- 2000 species – mostly freshwater
- Digestive tract with mouth and anus – mastax
- **Pseudocoelomate**
- Protonephridia with flame bulbs

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Reproduction

- **PARTHENOGENESIS** – unfertilized diploid eggs develop into females (all Class Bdelloidea rotifers are female)
- Unfertilized haploid eggs become degenerate males that live long enough to produce sperm, with no digestive tract
- Loss of sex reproduction is simple allelic Mendelian inheritance
- Sexual dimorphism

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



A ROTIFER



ROTIFERS



Rotifers have cilia on their head which can create a vortex to suck in food

Mag. 400X

ROTIFER EATING



LOPHOPHORATES

Phylum Bryozoa

Phylum Brachiopoda

Lophophore – ciliary feeding device that also functions in respiration

True coelom

Phylum **Bryozoa- “moss animals”**

- Small colonial aquatic animals
- Look (superficially) like plants
- About 4,000 species
- Animal (“zooid”) secretes and lives inside “zoecium” (of chitin, calcium carbonate, or gelatinous)
- Monoecious

Phylum **Brachiopoda – lamp shells**

- Marine with 2 shells
- Dorsal and ventral valve
- About 300 living species
- Well represented as fossils (12,000 species)
- Dioecious

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) A bryozoan



(b) A brachiopod, the northern lamp shell

BRACHIOPOD – “LAMP SHELL”



BRYOZOAN

BRYOZOAN COLONY



PHYLUM MOLLUSCA

Class Gastropoda

Class Bivalvia

Class Polyplacophora

Class Cephalopoda

Class Scaphopoda

PHYLUM MOLLUSCA



Table 11.1.

Copyright © McGraw-Hill Education. Permission required for reproduction or display.



TABLE 11.1
CLASSIFICATION OF THE MOLLUSCA

Phylum Mollusca (mol-lus′kah)
The coelomate animal phylum whose members possess a head-foot, visceral mass, mantle, and mantle cavity. Most molluscs also possess a radula and a calcareous shell. Nearly 100,000 species.

Class Caudofoveata (kaw′do-fo″ve-a″ta)
Wormlike molluscs with a cylindrical, shell-less body and scale-like, calcareous spicules; lack eyes, tentacles, statocysts, crystalline style, foot, and nephridia. Deep-water, marine burrowers. *Chaetoderma*. Approximately 150 species.

Class Solenogastres (so-len″ah-gas′trez)
Shell, mantle, and foot lacking; worm-like; ventral (pedal) groove; head poorly developed; surface dwellers on coral and other substrates. Marine. *Neomenia*. Approximately 250 species.

Class Polyplacophora (pol″e-pla-kof′o-rah)
Elongate, dorsoventrally flattened; head reduced in size; shell consisting of eight dorsal plates. Marine, on rocky intertidal substrates. *Chiton*.

Class Monoplacophora (mon″o-pla-kof′o-rah)
Molluscs with a single arched shell; foot broad and flat; certain structures serially repeated. Marine. *Neopilina*. Approximately 25 species.

Class Gastropoda (gas-trop′o-dah)
Shell, when present, usually coiled; body symmetry distorted by torsion; some monoecious species. Marine, freshwater, terrestrial. *Nerita, Orthaliculus, Helix*. More than 35,000 species.

Class Cephalopoda (sef″ah-lah″po′dah)
Foot modified into a circle of tentacles and a siphon; shell reduced or absent; head in line with the elongate visceral mass. Marine. *Octopus, Loligo, Sepia, Nautilus.* Approximately 550 species.

Class Bivalvia (bi″val′ve-ah)
Body enclosed in a shell consisting of two valves, hinged dorsally; no head or radula; wedge-shaped foot. Marine and freshwater. *Anodonta, Mytilus, Venus*. Approximately 30,000 species.

Class Scaphopoda (ska-fop′o-dah)
Body enclosed in a tubular shell that is open at both ends; tentacles used for deposit feeding; no head. Marine. *Dentalium*. More than 300 species.

This taxonomic listing reflects a phylogenetic sequence. The discussions that follow, however, begin with molluscs that are familiar to most students.

Copyright © 2016 McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.

MOLLUSCAN CHARACTERISTICS

1. Body of two parts: head-foot and **visceral mass**
2. **Mantle** that secretes a calcareous shell and covers the visceral mass
3. Mantle cavity functions in excretion, gas exchange, elimination of digestive wastes, and release of reproductive products.
4. Bilateral symmetry
5. **Trochophore** larvae, spiral cleavage, and schizocoelous coelom formation
6. Coelom reduced
7. **Open** circulatory system (except Cephalopoda)
8. **Radula** usually present

BODY ORGANIZATION

Head-foot

- Mouth
- Foot for attachment and locomotion

Visceral mass

- Organs of digestion, circulation, reproduction, excretion

Mantle

- Secretes shell
- Respiration

Mantle cavity

- Excretion, elimination of digestive wastes and reproductive products, feeding in bivalves, movement in cephalopods

Radula

- Rasping food

FIGURE 28.20



There are many species and variations of mollusks; this illustration shows the anatomy of an aquatic gastropod.

Molluscan body organization.



CLASS GASTROPODA

GASTROPODS: SNAILS, LIMPETS, SLUGS, WHELKS, CONCHES, PERIWINKLES, SEA SLUGS (NUDIBRANCHS), SEA HARES, AND SEA BUTTERFLIES



SEA HARE



NUDIBRANCHS

CLASS GASTROPODA

- *Torsion*
  - Developmental process that changes the relative position of the shell, digestive tract and anus, nerves that lie on both sides of the digestive tract, and the mantle cavity containing the gills



In **veliger** larva, anus is posterior; no mantle cavity is visible

Shell rotates to adult position; viscera rotates 90˚; mantle cavity visible

Anus and mantle continue to rotate; mantle cavity expands

Anus rotated 180˚; mantle epithelium reorganizes to enlarge cavity

Adult snail

SNAIL SHOWING TORSION



By User:Fir0002 - Own work, GFDL 1.2, https://commons.wikimedia.org/w/index.php?curid=5520991

CLASS GASTROPODA

Coiling

- Spiral winding of the shell and the visceral mass inside.
- Ancestors had a flat spiral shell – not as compact.
- Modern snail form spirals out to the side, but it's heavy, so shell shifted up higher.

Locomotion

- Flattened foot
- Cilia propel over mucous trail
- Muscular waves

CLASS GASTROPODA

© Clouds Hill Imaging/www.lastrefuge.co.uk

Feeding and digestion

- Most scrape algae and attached organisms
- Herbivores, predators, scavengers
- Oyster borer alternates rasping with chemical softening of the shell to bore a hole in an oyster





GASTROPOD FEEDING DIVERSITY

- Species of *Conus* deliver a *lethal sting to secure prey*
  - *Venom* is specific for the neuroreceptors of its preferred prey
  - Radula tooth modified into a hypodermic needle and this is attached to a venom gland
  - Tooth is barbed on the end of a proboscis and can be extended out of the head.

CONUS

OTHER BODY FUNCTIONS

Open circulatory system

- Blood bathes tissues in sinuses.
  - Transports nutrients, wastes and gases

Nervous system

- Six ganglia plus nerve cords

Sensory structures

- Eyes at base or end of tentacles
- Statocysts in foot



CLASS GASTROPODA FACTS

- Pulmonates (= with lungs)
  - Includes all land and most freshwater snails and slugs
  - Ancestral gill has been lost and the mantle wall (full of blood vessels) is now a lung
  - Air fills lung by contraction of mantle floor

Monoecious.
Slug sex video

By SeanMack - Own work, CC BY 3.0, https://commons.wikimedia.org/w/index.php?curid=2830365

DAVID ATTENBOROUGH SLUG SEX

SNAIL TENTACLES



Aquatic species have one pair of tentacles.

Landforms have two pair of tentacles and the posterior pair has eyes.





CLASS BIVALVIA

CLASS BIVALVIA

Class Bivalvia

Mussels, clams, scallops, oysters, and shipworms

Range in size from 1–2 mm in length to the giant South Pacific clams

Most are *sedentary filter feeders*

Bivalves *lack a head, radula, or other aspects of cephalization*

Most are marine



BIVALVE SHELL

2 shells or “valves”

- Valves are drawn together by strong *adductor muscles*

Umbo is the oldest part of the shell with growth occurring outward in rings

Pearls are produced when an irritant is lodged between the shell and mantle

- *Layers of nacre* are secreted around the foreign material

PEARLS

Figure 11.3 Molluscan shell and mantle.



© BIODIDAC - Fins éducatives seulement / Educational use only

GAS EXCHANGE, FILTER FEEDING, AND DIGESTION

Sedentary filter feeders (like clams)

- Loss of head and radula
- Cilia-covered gills in folded sheets
- Cilia create water currents into and through mantle cavity.
    - Food is trapped along gill surface and transported to food grooves and labial palps toward mouth.

Bivalve feeding. Blue arrows show path of water into and out of clam. Yellow arrows show path of food particles, which come in with the water current and get stuck on the gills.



OTHER BODY FUNCTIONS

Open circulatory system

- Mantle and gills oxygenate blood

Nephridia (liquid nitrogenous wastes)

- Below pericardial cavity

Nervous system

- Three pairs of interconnected ganglia
- Sensory receptors at mantle margin



© 2001 Sinauer Associates, Inc.



Livingstone, © BIODIDAC.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Shipworms



A



Larry S. Roberts

B



Larry S. Roberts





Shipworms - *Teredo*

REPRODUCTION AND DEVELOPMENT

Mostly dioecious

Gonads within visceral mass

Mostly external fertilization

Trochophore and veliger larval stages

Freshwater group has

- Parasitic larval stage (parasitic on fish gills), called a glochidium

Trochophore (a) and veliger (b) larval stages. Glochidia of an freshwater bivalve.







Trochophore larva



Livingstone, © BIODIDAC.



prehatch veliger

Glochidium Larva :- Found In **Mollusca**



Glochidium Larva

> **CATLIN SEAVIEW SURVEY** - Ribbon Reef No. 5 - 31/10/12

© Underwater Earth - Jayne Jenkins

Bivalve diversity. Rock scallop (*Hinnites giganteus*).



© Takako Uno

ROCK SCALLOP



Bivalve diversity. The geoduck



Digging deep

The geoduck lives a sedentary life beneath the seafloor. Adults dig to a depth of about 3 feet. If unearthed, they cannot rebury themselves.



Once situated, the geoduck pulls in water, filtering food, and expels waste through the siphon neck.

Young geoduck extends foot.

Then expands foot to anchor itself in the sand.

By contracting its foot muscles...

...the clam pulls its shell downward.

Adult geoduck burrowed and secure.



Geoduck - ***Panopea abrupta***

CLASS CEPHALOPODA

Squid, octopuses, cuttlefish, and nautiluses

Foot modified into circle of tentacles or arms and incorporated into siphon

Head in line with visceral mass

Muscular mantle

REPRESENTATIVE CEPHALOPODS

Squid and Cuttlefish (8 arms and 2 tentacles)

Chambered Nautilus (up to 90 tentacles)

Octopus and Vampire squid (8 arms)

Most complex brain of any invertebrate (memory and learning)

Head and foot completely merged, and the arms, tentacles and siphon are derived from the foot

Have beak and radula, some with poison salivary glands

No larval stage, semelparous (mate once and die)

Chromatophores and skin texture change

Eyes with cornea, lens, retina and iris

Internal structure of the squid

Squid (Giant)

Architeuthis dux



©Sheri Amsel

www.exploringnature.org

CLASS CEPHALOPODA

Shell

- Reduced or absent except in nautilus

Locomotion

- Jet propulsion using muscles of mantle compressing water within mantle cavity and siphon

Feeding and Digestion

- Predators
  - Tentacles, jaws, radula (beak)
  - Octopus tenting behavior
- Digestive tract muscular with large digestive glands

OTHER BODY FUNCTIONS

A. Closed circulatory system

B. Nervous system

   1. Large brains (wrap around their esophagus)
      a. Best learning ability for an invertebrate
   2. Complex sensory structures
      a. Eyes
      b. Statocysts

C. Chromatophores

   1. Color changes involved with courtship and other displays
   2. Cuttlefish colors

D. Ink glands

The cephalopod eye.



REPRODUCTION AND DEVELOPMENT

Dioecious

Male produces spermatophores

- Transfers to female’s mantle cavity using modified tentacle (hectocotylus)

Eggs deposited singly or in masses attached to substrate.

Eggs tended by parents.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Mating in cephalopods



CEPHALOPODS

- Include: squid, octopus, nautilus and cuttlefish

- Considered most derived and advanced of the protostomes

- Highly successful in oceans

- Structural complexity exceeds all other invertebrates

SQUID

- Most successful of the cephalopods
- Have 8 fully suckered arms and 2 longer tentacles with spatulate suckered tips (octopi only have 8)
- Reduced shell
- Siphon is used for locomotion and can be moved forward or back to change direction quickly (jet propelled)
- Developed a camera-like eye similar in function to vertebrate eyes (NOT homologous)

STRANGE SQUID EYES

Architeuthis clarkei: the giant squid

David Paul

OCTOPUS

COCONUT OCTOPUS

NEW SPECIES OF OCTOPUS



CUTE OCTOPUS



CHAMBERED NAUTILUS



A



B



C

A: Richard J. Neves

CUTTLEFISH



CLASS POLYPLACOPHORA

Chitons

Reduced head, flattened foot, shell consisting of eight dorsal valves, muscular mantle extends beyond margin of shell (figure 11.20)

Feed on attached algae

Ladderlike nervous system

Dioecious with external fertilization



CHITON VIDEO

CLASS SCAPHOPODA

Tusk shells

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

The scaphopodan *Dentalium*





ANNELIDS AND ALLIED TAXA

Venomous fireworm with warning coloration

CHAPTER OPENER 11

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Photo Gear#CRAB 02.tif

FIGURE 28.30


(a)
(b)
(c)

The (a) earthworm, (b) leech, and (c) featherduster are all annelids. (credit a: modification of work by S. Shepherd; credit b: modification of work by “Sarah G...”/Flickr; credit c: modification of work by Chris Gotschalk, NOAA)

Lophotrochozoa - continued

Members have either a **lophophore** (a crown of ciliated tentacles)

- Bryozoans
- Brachiopods
- Rotifers

Or have a **trochophore larval stage**

- **ANNELIDS**
- Molluscs
- (Platyhelminthes (flatworms))



Structure of ??? larva

PHYLUM ANNELIDA

Lophotrochozoans

Common ancestry

with Molluscs

Current classification

- Polychaetes
- Earthworms and leeches

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



WHAT’S IN AN ANNELID?

1. Body segmented, bilaterally symmetrical, and worm-like
2. Spiral cleavage, trochophore larvae
3. Paired epidermal setae (bristles of chitin)
4. Closed circulatory system (dorsal and ventral blood vessels)
5. Ventral nerve cord with ganglia
6. Nephridia for excretory (liquid wastes)
7. Respiration by diffusion with moist conditions

ANNELID DIVERSITY, THE SEGMENTED WORMS: ERRANT POLYCHAETE (LEFT), FEATHER-DUSTER WORM (ALSO A POLYCHAETE[MIDDLE]), LEECH (RIGHT), TUBIFEX (BOTTOM)



DIVIDING THE BODY

Annelids exhibit segmentation

Bodies composed of serially repeated units

- Each unit contains components of most organ systems



Coelom

- Septa are membranes of epithelial tissue that separate segments of annelid
- Gut and longitudinal blood vessels extend through the septa



Copyright © McGraw-Hill Education. Permission required for reproduction or display.



Evolution of segmentation allowed much *greater complexity in structure and function*

- Increased burrowing efficiency by permitting independent movement of segments (hydrostatic skeleton)
- Evolution of a more sophisticated nervous system
- Provided a safety factor
    - If one segment failed, others could still function

CHARACTERISTICS

- *Trochophore* is the ancestral ciliated larval form
- Segmented worms living in marine, freshwater, and moist terrestrial habitats
- Phylum Annelida includes polychaetes (marine bristle worms), leeches, and earthworms





TROCHOPHORE LARVAE DEVELOPMENT



PHYLUM ANNELIDA

- About 15,000 species
- 2/3 are the more obscure marine worms.
- *Setae*
  - Tiny chitinous bristles called setae (Absent in leeches)
  - Short setae anchor segments while moving in the soil for earthworms
    - Prevent it from slipping backward, help prevent a worm from being pulled or washed out of its home
  - Long setae help aquatic worms swim or crawl
  - Hold marine worms in their burrows





Karen Osborn, Smithsonian In


Longitudinal muscle
Formative cell
Circular muscle
Seta
Epidermis
Retractor muscle
Cuticle
Peritoneum
Protractor muscle

Polychaete locomotion.



PHYLUM ANNELIDA

Phylogeny

Traditionally, annelids are divided among *3 classes*

- Class Polychaeta
- Class Oligochaeta
- Class Hirudinida

Now, 2 classes:

- Class Polychaeta: Mobile Polychaetes, sedentary Polychaetes (tube worms)

- Class Oligochaeta: Oligochaetes (earthworms), Hirudinida (leeches)

PHYLUM ANNELIDA

Polychaetes

- Primarily marine and usually benthic (in the sand)
- On ocean floor, under rocks & shells, crevices of coral reefs, burrows.
- Parapodia usually the **chief respiratory organ** for polychaetes, although marine forms may also have gills



Earthworms and leeches

- Freshwaters, or terrestrial soils
- Many leeches are bloodsuckers,
  others are carnivores (predators)

PHYLUM ANNELIDA: EARTHWORM

Body Plan

Body Wall

- Head is composed of
    - *Prostomium and mouth*
- Surface is covered with an epidermis and a thin outer layer of non-chitinous cuticle
- Strong *circular and longitudinal muscles* underlie the body wall, for swimming, crawling and burrowing.



PHYLUM ANNELIDA

Coelom

- Forms by *schizocoely* (protostome development)



Ivy Livingstone © BIODIDAC

Guy/97

PHYLUM ANNELIDA

Hydrostatic Skeleton

- Except in leeches, coelom is *filled with fluid and serves as a hydrostatic skeleton*
- Fluid volume *remains constant*
- Contraction of *longitudinal muscles*
    - Causes body to shorten and expand
- Contraction of *circular muscles*
    - Causes body to narrow and lengthen
- Swimming annelids use undulatory movements

OLIGOCHAETES (EARTHWORMS)

Nutrition

- *Scavengers,* feeding on decayed organic matter and bits of vegetation
- Food moistened by mouth and drawn in by muscular pharynx
- Digestive tract is unsegmented and extends the length of the earthworm **(complete digestive tract)**


To Part C below
Prostomium
Seminal receptacle openings
♀
♂
Genital openings
Clitellum
Setae
Anus
A
Pharynx
Lateral nerve
Second left aortic arch
Seminal vesicles
Esophagus
Epidermis
Gizzard
Circular muscle
Longitudinal muscle
Dorsal vessel
Brain
Buccal cavity
Prostomium
Crop
Intestine
Mouth
Testes
Ovary
B
Left circumpharyngeal connective
Seminal receptacle
Sperm funnel
Egg funnel and oviduct
Nerve cord
Subneural vessel
Nephridium
Ventral vessel
Circular muscle
Epidermis
Cuticle
Dorsal vessel
Longitudinal muscle
Peritoneum
Typhlosole
Intestinal lumen
Chloragogen cells
Intestinal epithelium
Setae
Setal retractor muscle
Nephridium
C
Lateroneural vessel
Ventral vessel
Subneural vessel
Ventral nerve cord
Mucous gland cell
Pores of gland
Sensory (receptor) cell of sense organ
Epithelial cell
Cuticle
Photoreceptor cell
Sensory fibers
D

OLIGOCHAETES

- Digestion and absorption occur in intestine
- *Typhlosole in intestines* increases surface area for nutrient absorption



Figure 12.18 Earthworm cross section.

Copyright © McGraw-Hill Education. Permission required for reproduction or display.



Copyright © 2016 McGraw-Hill Education.
All rights reserved. No reproduction or distri

OLIGOCHAETES

Circulation and Respiration

- *Coelomic fluid and circulatory system* transport food, wastes, and respiratory gases
- Blood circulates in a *closed system* with five main trunks running lengthwise in the body
- *Dorsal vessel is the main pumping organ*



Excretion

Most body segments have a *pair of nephridia*

Ends at a bladder that empties outside at *nephridiopore*



Human nephron



OLIGOCHAETES

Nervous System and Sense Organs

- Pair of cerebral ganglia connect around the pharynx to the ganglia of the ventral nerve cord to form the brain
- Fused ganglia in each segment
- One or more *giant axons* located in ventral nerve cord increase rate of conduction



OLIGOCHAETES

- *Earthworms are hermaphroditic*
- Mate at night during warm, moist weather
- Aligning in *opposite directions,* ventral surfaces together
- Mucus secreted by *clitellum* holds worms together
- After mutual copulation, each worm secretes a mucus tube and chitinous band to form a *cocoon*
- Cocoon passes forward and *eggs, albumin, and sperm are poured into it*
- *Fertilization* occurs in cocoon-Cocoon slides off head end and closes
- *Young worms* emerge from cocoon

EARTHWORM MATING



Any two earthworms can mate by joining together and fertilizing each other's eggs.

Later, a tube of mucus secreted by the clitellum forms an egg case, which slips off the worm's body.

Young worms hatch out into the soil.

G

© G.L.Twiest/Visuals Unlimited



Giant earthworm - Australia

Diversity

Largest group of annelids

More than 10,000 species, *mostly marine*

Vary from 5 to 10 cm long

Some live in crevices, others inhabit tubes

Sedentary polychaetes mainly tube-living, eat organic particles floating in sea water

Predatory or scavenging polychaetes may be free-moving, burrowing, or crawling

Important part of *marine food chains*

526 × 350 - australianmuseum.net.au

ERRANT POLYCHAETE



BOBBIT WORM – CAN GET TO 20 FT LONG

Copyright 2009 photolibrary.com



Late stage polychaete trocophore larva

POLYCHAETES

Well-differentiated head with sense organs

Prostomium (1st segment) often bears eyes, tentacles, and sensory palps

- Surrounds mouth, which may have setae, and/or chitinous jaws



www.bumblebee.org



POLYCHAETES

Paired appendages called *parapodia* on most segments

Many setae arranged in bundles on parapodia

- Parapodia used for crawling, swimming, or anchoring in tubes
- Usually the chief *respiratory organ* although the worm may also possess gills



POLYCHAETES

Tubeworms

- Tube-dwellers
- May line their burrows with mucus
- Use cilia or mucus to obtain food

Fanworms

- Unfurl tentacular crowns to feed
- Food moved from fan to mouth by ciliary action in a sac to help build tube
- Store sand grains


Atoke
Epitoke



Arenicola - lugworm

The lugworm *Arenicola*



Lugworm casts



By Original uploader was Nveitch at en.wikipedia - Transferred from en.wikipedia, CC BY-SA 3.0, https://commons.wikimedia.org/w/index.php?curid=4081034


Ollie and Quentin
by Piers Baker
www.ollieandquentin.com
©2011 Piers Baker. Distributed by King Features Syndicate
03-27
?
In case of memory loss
Ollie
(seagull)
Quentin
(lugworm)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

A sedentary polychaete (*Chaetopterus*)



FAMILY HIRUDINIDAE

Leeches

- Anterior and posterior suckers
- *Most freshwater,* few marine or live in moist terrestrial environments
- *Colors:* black, brown, red, and olive green
- Lack distinct coelomic compartments
    - No septa
- *No setae*



Photograph by T. Branning

FAMILY HIRUDINIDAE

Nutrition



- Not all are parasites, *many are predaceous,* feeding on small invertebrates
- Freshwater leeches have a proboscis for ingesting small invertebrates as well as to suck blood or pharynx with 3 jaws armed with teeth
- Most are *fluid feeders* that prefer tissue fluids and blood pumped from open wounds
- anticoagulant and narcotic in their saliva
- Gut specialized for storage and digestion of large quantities of blood-

Figure 12.20 Internal Structure of a Leech.

Copyright © McGraw-Hill Education. Permission required for reproduction or display.



Copyright © 2016 McGraw-Hill Education.
All rights reserved. No reproduction or distri

Leech locomotion.



LEECH
MOVEMENT

Cleveland P. Hickman, Jr.

MEDICINAL LEECHES

Medical Science Opens Up a Can of Worms With Leech Therapy : Health: Doctors use the parasite to promote blood circulation in patients who have had severed fingers, eyelids, ears or other extremities surgically reattached.



[Biopharm]

[Biopharm web page]



About us

Biopharm is an international company based in Hendy, South Wales, UK. Established originally in 1812, it moved to its current base in 1984. Biopharm is the first leech farm of its kind and we are innovators in the cultivation of the medicinal leech, currently producing the majority of leeches used in modern medicine worldwide.

Biopharm is the pioneer in medical leech provision and has nurtured the renaissance of leeches in modern surgery. Leeches today are used in plastic and reconstructive surgery worldwide, as well as successfully ongoing research into relieving symptoms of osteoarthritis. Their use in veterinary surgery, especially in polycythaemia vera has been particularly successful.

Our leeches are maintained in as sterile and pure environment as is possible; we

Dan Winston recommends Biopharm Leeches Ltd.
Applicator at sealant applicator · 137 followers · 22 July

Cant recommend Biopharm enough, product is amazing
And sales excellent arrive in great condition in quick time, usually order before 2 one day and arrive same n...

See more

WHAT’S IN AN ANNELID? SUMMARY

1. Body segmented, bilaterally symmetrical, and worm-like
2. Spiral cleavage (Protostome development), trochophore larvae
3. Paired epidermal setae (bristles of chitin)
4. Closed circulatory system with pseudohearts and dorsal & ventral blood vessels
5. Ventral nerve cord with ganglia
6. Nephridia for excretory (liquid wastes)
7. Respiration by diffusion with moist conditions



v · t · e **Extant phyla of kingdom Animalia** [hide]

Domain: Archaea · Bacteria · Eukaryota · (Supergroup: Plant · Hacrobia · Heterokont · Alveolata · Rhizaria · Excavata · Amoebozoa · Opisthokonta (**Animal** · Fungi))

Basal / incertae sedis								**Porifera (sponges)** · Ctenophora (comb jellies) · Placozoa (*Trichoplax*) · Mesozoans (Orthonectida · Dicyemida or Rhombozoa)
Planulozoa								**Cnidaria (jellyfish and relatives)**
	Bilateria	**Xenacoelomorpha**						Xenoturbellida *(Xenoturbella)* · Acoelomorpha (Acoels · Nemertodermatids)
		Nephrozoa	**Deuterostomia**					**Chordata (lancelets · tunicates · craniates / vertebrates)**
				Ambulacraria				**Echinodermata (starfish and relatives)** · Hemichordata (Acorn worms · Pterobranchs)
			Protostomia	*Basal / incertae sedis*				Chaetognatha (arrow worms)
				Ecdysozoa	**Scalidophora**			Kinorhyncha (mud dragons) · Priapulida (penis worms) · Loricifera
					Nematoida			**Nematoda (roundworms)** · Nematomorpha (horsehair worms)
					Panarthropoda			**Arthropoda (arthropods)** · Tardigrada (waterbears) · Onychophora (velvet worms)
				Spiralia	**Gnathifera**			Gnathostomulida (jaw worms) · Micrognathozoa *(Limnognathia)* · Syndermata (Rotifera · Acanthocephala) · Cycliophora *(Symbion)*
					Platytrochozoa	**Rouphozoa**		**Platyhelminthes (flatworms)** · Gastrotricha (hairybacks)
						Lophotrochozoa	**Lophophorata**	**Bryozoa (moss animals)** · Brachiozoa (Brachiopoda or lamp shells · Phoronida or horseshoe worms)
							Other clades	**Annelida (ringed worms)** · **Mollusca (molluscs)** · Nemertea (ribbon worms) · Entoprocta (Kamptozoa)

Major groups within phyla	**Sponges** (Calcareous · Hexactinellid · Demosponge · Homoscleromorpha) · **Cnidarians** (Anthozoa inc. corals · Medusozoa inc. jellyfish · Myxozoa) · **Vertebrates** (Jawless fish · Cartilaginous fish · Bony fish · Amphibians · Reptiles · Birds · Mammals) · **Echinoderms** (Sea lilies · Asterozoa inc. starfish · Echinozoa) · **Nematodes** (Chromadorea · Enoplea · Secernentea) · **Arthropods** (Chelicerates/Arachnids · Myriapods · Crustaceans · Hexapods/Insects) · **Platyhelminths** (Turbellaria · Trematoda · Monogenea · Cestoda) · **Annelids** (Polychaetes · Clitellata · Echiura) · **Molluscs** (Gastropods · Cephalopods · Bivalves · Chitons · Tusk shells)

Phyla with ≥5000 extant species **bolded** · See also: Parazoa and Eumetazoa; Radiata; Platyzoa; Monoblastozoa (*nomen dubium*)

SUPERPHYLUM LOPHOTROCHOZOA
FLATWORMS
PHYLUM PLATYHELMINTHES

Lophotrochozoa

Members have either a **lophophore** (a crown of ciliated tentacles)

- Bryozoans
- Brachiopods
- Rotifers

Or have a **trochophore larval stage**

- Annelids
- Molluscs

- Platyhelminthes (flatworms) share 18sRNA with the lophophorates, and some have cilia in the pharynx for feeding


Multicellular
Porifera
Cnidaria
Loph
Platyhelminthes
Annelida
Protostome development
Mollusca
True tissues
Ecdys
Nematoda
Arthropoda
Notochord
Deuterostome development
Chordata
Bilateral symmetry
Echinodermata

LOPHOTROCHOZOA





Trochophore larva and lophophore ciliated feeding tentacles.

Flatworms share 18sRNA with lophophorates and some have feeding cilia.

CHARACTERISTICS OF LOPHOTROCHOZOANS

1. Triploblastic
   a) Mesoderm between endoderm and ectoderm
2. Protostomes
   a) Schizocoely
   b) Mouth first
3. Bilateral symmetry
4. Cephalization
5. Trochophore larvae or lophophore feeding structure

GENERAL FEATURES

Animals that actively seek food, shelter, home sites, and mates require a different set of strategies and body organization than sessile organisms

Two major evolutionary advances in animal evolution:

- *Cephalization*
    - Concentrating sense organs in the head region
- *Primary bilateral symmetry*
    - Body can be divided along only 1 plane of symmetry to yield 2 mirror images of each other
    - “Worm” - an elongated body form with anterior (head), posterior (head), dorsal, and ventral sides

FLATWORMS

These are the simplest animals with *primary bilateral symmetry and cephalization*

Triploblastic and acoelomate, with ectoderm, mesoderm, endoderm

Some are highly specialized *parasites, others free-living and predatory*

FIGURE 28.14 FLATWORM GROUPS



(a) Class Turbellaria

(b) Class Monogenea



(c) Class Trematoda

(d) Class Cestoda

Phylum Platyhelminthes is divided into four classes.

(a) (Class) Turbellaria includes the Bedford’s flatworm (*Pseudobiceros bedfordi*), which is about 8–10 cm in length.

(b) The parasitic class Monogenea includes *Dactylogyrus* spp, commonly called a gill fluke, is about 0.2 mm in length and has two anchors, indicated by arrows, that it uses to latch onto the gills of host fish.

(c) The Trematoda class includes *Fascioloides magna* (right) and *Fasciaola hepatica* (two specimens of left, also known as the common liver fluke).

(d) Class Cestoda includes tapeworms such as this *Taenia saginata.* which infects both cattle and humans, and can reach 4–10 meters in length. (credit a: modification of work by Jan Derk; credit d: modification of work by CDC)

PHYLUM PLATYHELMINTHES



	Cnidarians and Ctenophores[4]	Platyhelminthes (including flatworms)[3][6]	More "advanced" bilaterians[8]
Bilateral symmetry	No	Yes	
Number of main cell layers	Two, with jelly-like layer between them	Three	
Distinct brain	No	Yes	
Specialized digestive system	No	Yes	
Specialized excretory system	No	Yes	
Body cavity containing internal organs	No		Yes
Specialized circulatory and respiratory organs	No		Yes

PHYLUM PLATYHELMINTHES

- Primitive NERVOUS and EXCRETORY (protonephridia and FLAME CELLS) systems: no circulatory, respiratory or skeletal systems.
- Highly developed reproductive system in parasitic forms: genital openings often on the ventral surface. Most are MONOECIOUS, a few are DIOECIOUS.
- Well developed MUSCULAR system

FIGURE 28.13 DIGESTIVE, EXCRETORY, NERVOUS



The planarian is a flatworm that has a gastrovascular cavity with one opening that serves as both mouth and anus. The “excretory” system is made up of tubules connected to excretory pores on both sides of the body. The nervous system is composed of two nerve cords running the length of the body, with cerebral ganglia and eyespots at the anterior (head) end.

SENSE ORGANS

Active locomotion favored *cephalization* and evolution of *sense organs*

Ocelli (light-sensitive eyespots)

- Present in turbellarians and larval trematodes

Touch sensitive and chemical sensitive cells

- Abundant, especially in the ear-shaped *auricles*

Sensory nerve endings found in

- Oral suckers and genital pores of parasitic groups

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

PHYLUM PLATYHELMINTHES

Characteristics

- Vary from a millimeter to many meters in length
- Many free-living marine forms are colorful and predatory

Marine flatworms



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.



Marine Flatworms-Facts-Photographs an...
seaunseen.com

Marine Flatworms-Facts-Photographs and ...
seaunseen.com

Marine flatworms: The butterflies of ...
africageographic.com

Diving in Bali. Interesting facts ...
diveguideinbali.com

Marine flatworms: The butterflies of ...
africageographic.com

Photo #86, Pseudoceros cf. dimidiatus
rzuser.uni-heidelberg.de

The Sea Slug Forum - Flatw...
seaslugforum.net

31 best Marine Flatworms images ...
pinterest.com

Phylum Platyhelminthes, Flatworms
marinelifephotography.com

marine worms: Polyclad and Acoela ...
starfish.ch

Marine Flatworms of the World! Photo ...
rzuser.uni-heidelberg.de

World's Weirdest: Flatworm Penis Fencing
video.nationalgeographic.com

Marine Flatworm; Acanthozoon specie...
chaloklum-diving.com

flatworms - Phylum Platyhelminthes
ryanphotographic.com

Flatworm marine.jpg - Wikimedia Co...
commons.wikimedia.org

Marine flatworms
slideshare.net

Marine Flatworms of the World ...
rzuser.uni-heidelberg.de

Flatworms: Looks can be deceiving ...
blog.nus.edu.sg

Marine Flatworms-Facts-Photographs and ...
seaunseen.com

Ocean Beauties: Marine Flatworms
leisurepro.com

Marine Flatworms of the World! Phot...
rzuser.uni-heidelberg.de

Marine Flatworms of the World! Photo ...
rzuser.uni-heidelberg.de

Diving in Bali. Interesting facts ...
diveguideinbali.com

Marine flatworms: The butterflies of ...
africageographic.com

GENERAL FEATURES

Triploblastic animals can be acoelomate, pseudocoelomate, or true coelomate

Acoelomates (flatworms)

- Typical acoelomates have only one internal space, the digestive cavity
  - Region between the epidermis and digestive cavity is filled mesodermal tissue



Planaria cross section, pharynx region

(CLASS) TURBELLARIA

- Mostly free-living forms
- Most are bottom dwellers in marine or freshwater
- Freshwater planarians
  - Found in streams, pools, and hot springs
- Terrestrial flatworms limited to moist places

Arrowhead flatworm



MARINE FLATWORMS

They are carnivorous and eat things like small fish, tunicates, small crustaceans, worms, clams, and molluscs. Some even eat algae and some feed on dead organisms instead of alive ones. Most species of marine flatworms have specialized diets. In addition, some flatworm species form a symbiotic relationship with algae, in which the algae live inside the worm and photosynthesize, producing sugar for the flatworm to use as energy.

https://www.scuba.com/blog/explore-the-blue/marine-flatworms/



ARROWHEAD FLATWORM. (WE HAVE THESE IN FRESNO!)



CLASS TURBELLARIA MOVEMENT

PHYLUM PLATYHELMINTHES



(Class) Turbellaria

- Range from 5 mm to 50 cm long
  - Ventral mouth and pharynx that is half-way down the body
  - Some have simple gut or no gut and a simple pharynx with cilia
  - Others have branched gut/intestine with an eversible pharynx
  - Flame cells

No respiratory organs or true excretory organs. How do they do it?


Intestine
Pharynx
Mouth
A
Mouth
Pharynx
Intestine
B

CLASS TURBELLARIA

Epidermis and Muscles

- Most have *cilia* on ventral side, attached to epidermis
  - Glide over a *slime track* secreted by adhesive glands
- Rhythmical muscular waves pass backward from the head



BIODIDAC ©, J. Houseman

FEEDING

- Planaria
  - Carnivorous and detect food by *chemoreceptors*
  - Food trapped in mucous secretions from glands
  - Wrap themselves around prey
  - Extend the proboscis to suck up bits of food

PLANARIA EATING

REPRODUCTION AND REGENERATION

Fission

- Many turbellarians constrict behind the pharynx and separate into two animals
- Each half regenerates the missing parts
  - Provides for *rapid population growth*

Regeneration

- If the head and tail are cut off
  - Each end grows the missing part; it *retains polarity*
- *Extract of heads* added to a culture of headless worms *prevents regeneration*



SEXUAL REPRODUCTION IN PLANARIA



These monecious flatworms reproduce sexually by “penis-fencing”. Each tries to stab the other with its penises. The successful one releases sperm into the “loser”, who then must develop eggs and care for them internally until the eggs hatch.

CLASS MONOGENEA









Gyrodactylus sp



D. PARADOXUM, A CLASS MONOGENEA FLATWORM



Diplozoon paradoxum are hermaphroditic flatworms that live on gills of freshwater fishes. Males and females meet each other during their juvenile stage, and their bodies completely fuse together, whereupon they remain faithful until death," Dr. Barash said. Once the two worms have fused together cross-fertilization can be permitted. "That's the only species I know of in which there seems to be 100 percent monogamy."

Class Trematoda

- All trematodes are parasites called “flukes”
- Most adults are *endoparasites* of vertebrates

inside

The major group of Trematodes have at 2 or more hosts in their life cycle. (Digenea)

Most complex life cycles in the animal kingdom.

CLASS TREMATODA (FLUKES)

- *Adaptations for parasitism include:*
  - Penetration glands (to get into hosts)
  - Glands also secrete a cuticle to prevent digestion by host.
  - Hooks and suckers for adhesion
  - Increased reproductive capacity (body mostly reproductive organs)

Nutrition and Digestion

Mouth of *trematodes* (flukes)

- Opens near the anterior end ("head" end)
- Pharynx is not extensible



CLASS TREMATODA (FLUKES)

- Flukes feed on host cells, cellular debris, and body fluids
- Undigested food regurgitated out of its pharynx

SCIENCEPHOTOLIBRARY

DIGENEAN LIFE HISTORY STAGES

- ***MIRACIDIUM (CILIATED) – in mollusc or annelid***
- ***SPOROCYST***
- ***REDIA***
- ***CERCARIA***
- ***METACERCARIA***
- ***ADULT***



CLASS TREMATODA

Sheep Liver Fluke

Adult fluke lives in bile passageways in the liver of sheep and other ruminants.

Eggs are pass out in feces

Eggs hatch and miracidium larvae *penetrate snails* to become “sporocysts” (clones of the larvae)

Next stage (cercaria larvae) hatch and swim to vegetation.

- Encyst on vegetation and await being eaten by sheep

When eaten, develop into young flukes

CLASS TREMATODA (FLUKES)

Clonorchis sinensis: Human Liver Fluke

Most important human liver fluke

Common in China, Japan, and Southeast Asia

Also infects *cats, dogs, and pigs*

Adult fluke is 10–20 mm long with an oral and ventral sucker

Digestive system includes pharynx, esophagus, and two long intestinal ceca


fineart america



CLASS TREMATODA (FLUKES)



Clonorchis Life Cycle

- Adults live in *bile passageways* of humans and other fish-eating mammals
- Eggs are shed into water with feces
- The eggs hatch, larva swims, and burrows into snail. Transforms into a *sporocyst*
- Sporocyst clonally produces one generation of *clonal individuals (redia)*

CLASS TREMATODA



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Life cycle of *Clonorchis*

Metacercarial cysts in fish muscle
Liver
Bile duct
Adult fluke
Egg containing miracidium
Miracidium hatches after being eaten by snail
Sporocyst
Redia
Cercaria

CLASS TREMATODA

- Clones pass into the snail liver
  - Develop into *tadpole-like swimming cercaria larvae*
- Swimming larvae escape into water
  - Make contact with a fish
  - *Bore into fish muscles or under scales*
  - Shed tail and *encysts*
- A mammal eats *raw fish*
  - Cyst dissolves and flukes migrate up bile duct
- Heavy infection can *destroy the liver* and *result in death*
- Control of parasites
  - Destroy snails and thoroughly cook fish

CLASS TREMATODA

Schistosoma: Blood Flukes

Over 200 million people infested with *schistosomiasis*

- Common in Africa, South America, West Indies, and the Middle and Far East

Sexes are *separate*

3 species account for most human schistosomiasis:

Human to snail to human, penetrate skin

Live in abdominal veins



① Mature flukes live in the blood vessels of the human intestine. A female fluke fits into a groove running the length of the larger male's body, as shown in the LM at right.

② Blood flukes reproduce sexually in the human host. The fertilized eggs exit the host in feces.

③ If the human feces reach a pond or other source of water, the eggs develop into ciliated larvae. These larvae infect snails, the intermediate host.

④ Asexual reproduction within a snail results in another type of motile larva, which escapes from the snail host.

⑤ These larvae penetrate the skin and blood vessels of humans working in fields irrigated with water contaminated with fluke larvae.

SCHISTOSOMIASIS



CLASS TREMATODA

Other Trematodes

Leucochloridium

- Produces remarkably colorful sporocysts in snails' heads
- Tentacles pulse to attracts birds to eat snails and continue the life cycle



© - josef hlasek
www.hlasek.com
Succinea_putris
Leucochloridium macrostomum aa7261

ZOMBIE SNAILS



Disease	Vector	Population at risk (million)^1	Number of people currently infected or new cases per year	Present distribution	Likelihood of altered distribution
Malaria	Mosquito	2,400^2	300-500 million	Tropics and Subtropics	🟥
Schistosomiasis	Water snail	600	200 million	Tropics and Subtropics	🟨
Lymphatic Filariasis	Mosquito	1 094^3	117 million	Tropics and Subtropics	🟩
African Trypanosomiasis (Sleeping sickness)	Tsetse fly	55^4	250 000 to 300 000 cases per year	Tropical Africa	🟩
Dracunculiasis (Guinea worm)	Crustacean (Copepod)	100^5	100 000 per year	South Asia, Arabian Peninsula, Central-West Africa	☐
Leishmaniasis	Phlebotomine sand fly	350	12 million infected, 500 000 new cases per year^6	Asia, Southern Europe Africa, Americas	🟩
Onchocerciasis (River blindness)	Black fly	123	17.5 million	Africa, Latin America	🟨
American Trypanosomiasis (Chagas disease)	Triatomine bug	100^7	18 million	Central and South America	🟩
Dengue	Mosquito	1,800	10-30 million per year	All Tropical countries	🟨
Yellow Fever	Mosquito	450	more than 5 000 cases per year	Tropical South America Africa	🟩

1. Top three entries are population-prorated projections, based on 1989 estimates.
2. WHO, 1994.
3. Michael and Bundy, 1995.
4. WHO, 1994.
5. Ranque, personal communication.
6. Annual incidence of visceral leishmaniasis; annual incidence of cutaneous leishmaniasis is 1-1.5 million cases/yr (PAHO, 1994).
7. WHO, 1995.

🟥 Highly likely 🟨 Very likely 🟩 Likely ☐ Unknown

GRID Arendal UNEP
GRAPHIC DESIGN : PHILIPPE REKACEWICZ

Source: Climate change 1995, Impacts, adaptations and mitigation of climate change: scientific-technical analyses, contribution of working group 2 to the second assessment report of the intergovernmental panel on climate change, UNEP and WMO, Cambridge press university, 1996.

Protozoan

Trematode

Nematode

Protozoan

Nematode

Protozoan

Nematode

Protozoan

Virus

Virus



M PLATYHELMINTHES

Class Cestoda

- Tapeworms have long flat bodies with *scolex*
  - with suckers and hooks
- Scolex is followed by a series of reproductive units or *proglottids*
- No digestive system
  - Rely on the host’s digestive tract
  - Absorb digested nutrients
- Lack sensory organs



TAPEWORMS

FIGURE 28.15


4 Embryos develop into larvae in muscle.
5 Humans are infected by ingesting raw or undercooked meat.
Embryos penetrate the intestinal wall and circulate to musculature.
3
T. saginata
T. solium
6 Scolex attaches to intestine.
2
Cattle (T. saginata) and pigs (T. solium) become infected by ingesting vegetation contaminated by eggs or proglottids.
7 Proglottids form.
T. saginata
T. solium
1
Eggs and proglottids in feces are passed into the environment.

Tapeworm (*Taenia* spp.) infections occur when humans consume raw or undercooked infected meat. (credit: modification of work by CDC)


suckers
hooks


gravid proglottids (note uterus and ova)
mature proglottids
immature proglottids
scolex


Proglottids with reproductive structures
100 µm
Hooks
Scolex
Sucker

CLASS CESTODA

- Nearly all cestodes require two hosts
  - Adult is parasitic in the *digestive tract of the vertebrate*
- Over 1000 species of tapeworms known, infecting almost all vertebrates
- Most tapeworms *do little harm to host*

DOG FLEA TAPEWORM



DPDx

Dipylidium caninum

CDC



1 🔬 Gravid proglottids are passed intact in the feces or emerge from perianal region of either animal or human hosts.

2 The proglottids disintegrate and release the egg packets. 🔬

3 Larvae of the typical flea intermediate host ingests egg packets. Oncospheres hatch and develop into cysticercoids.

4 Flea larvae mature into adult fleas, which continue to harbor infective cysticercoids.

5 ☣ Definitive host is infected by ingesting fleas containing cysticercoids.

6 Scolex

7 Cats and dogs can harbor fleas infectious to humans.

Oncosphere
Cysticercoid
Incidental human transmission via ingestion of infected fleas.
Adult in small intestine

☣ Infective stage

🔬 Diagnostic stage

CLASS CESTODA

Beef Tapeworm

- Lives as an adult in the digestive tract of humans
- Juvenile form found in intermuscular tissue of cattle
- Mature adults can reach over 10 meters (32.5 feet) in length with over 2000 proglottids




Cysticercus eaten by human in rare beef
Evaginated cysticercus in upper intestine
Undercooked meat with living cysticercus
Yolk gland
Excretory canal
Nerve cord
Testes
Uterus
Sperm duct
Sperm
Genital pore
Vagina
Ovary
Gravid proglottid
Invaginated cysticercus
Shelled larva (in feces)
Cysts in muscle ("measly beef")
Grass, contaminated with eggs, ingested by cow



© Cabisco/Visuals Unlimited

CLASS CESTODA

Pork Tapeworm

Adults live in small intestine of humans

Juveniles live in muscles of pigs

Cysticercosis commonly occurs in eyes or the brain: blindness, epilepsy or death, if you swallow the eggs directly, rather than the cysts (oncospheres).

Infection can be avoided by eating thoroughly cooked pork


Oncospheres develop into cysticerci in muscle
Humans infected by ingesting raw or undercooked infected meat
Oncospheres hatch, penetrate intestinal wall, and circulate to musculature
3
T. saginata
T. solium
5
Scolex attaches to intestine
2
Cattle (T. saginata) and pigs (T. solium) become infected by ingesting vegetation contaminated by eggs or gravid proglottids
6
Adults in small intestine
T. saginata
T. solium
d
1
Eggs or gravid proglottids in feces and passed into environment
i = Infective Stage
d = Diagnostic Stage

Taenia solium, the pork tapeworm

Taenia solium





Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Cerebral cysticercosis

Larry S. Roberts

Cerebral cysticercosis, (neurocysticercosis) caused by T.

CLASS CESTODA

Fish Tapeworm

- Adults found in intestines of humans, dogs, cats and other mammals
- Immature stages found in crustaceans and fish
- Largest parasite of humans, usually about 30 feet, but possibly reaching up to 20 meters (65 feet) in length

RADIATA:
PHYLUM CNIDARIA - JELLIES, CORALS, AND THEIR RELATIVES
PHYLUM CTENOPHORA

REVIEW OF ANIMAL PHYLOGENY



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

CNIDARIANS: HYDROZOANS (TOP LEFT), JELLY (TOP RIGHT), SEA ANEMONE (BOTTOM LEFT), CORAL POLYPS (BOTTOM RIGHT)



PHYLUM CNIDARIA

Four classes of Cnidaria

- Hydrozoa
- Scyphozoa
- Cubozoa
- Anthozoa



PHYLUM CNIDARIA CHARACTERISTICS





- All are *aquatic* and mostly *marine*
- *Radial Symmetry*
  - No head: oral/aboral
- *Two body types*
  - Free-swimming **medusa**
  - Sessile **polyps**
- *Diploblastic* (2 body tissue layers)
  - Epidermis and gastrodermis
  - *Mesoglea="jelly":* matrix that lies between epidermis and gastrodermis
    - Made of mostly water with collagen protein and some other proteins and carbohydrates
- *Cnidocytes* contain **nematocysts**
  - Hairlike trigger – cnidocil
  - Some are sticky, others sting
- *Simple* muscles and nerves
  - Not true muscles (not of mesodermal origin)
- *Reproduction* both asexual and sexual (note **PLANULA** larvae)

First animal clade with true tissues

Gastrovascular cavity for *extracellular* digestion

- ☐ Allows ingestion of larger food particles
- ☐ Advance over sponge’s *intracellular* digestion

True nerve cells arranged in **nerve net**

- ☐ No central control organ – i.e. no cerebral ganglion or brain

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



The mesoglea is thin in polyps and thick in medusae.

POLYP AND MEDUSA FORMS OF CNIDARIANS





(a) Sea anemone: a polyp



(b) Jelly: a medusa

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

PHYLUM CNIDARIA

- *Incomplete gut:* gastrovascular cavity (mouth but no anus)
- *Tentacles* usually encircle mouth or oral region
- *Muscular contractions* by specialized epithelial cells





PHYLUM CNIDARIA

Nerve net

- Simple nervous system

Cnidaria: hydra (also jellyfish, corals, and anemones)



Copyright © 2002, Elsevier Science (USA). All rights reserved.

Nerve net: diffusely distributed network of neurons

No excretory or respiratory system

SHAPE OF LIFE

PHYLUM CNIDARIA

- Many cnidarians are *effective predators*
- Possible due to presence of a unique cell type, *the cnidocyte*

- *Cnidoctyes*
  - Produces one of over 20 types of stinging organelles: **nematocysts**



PHYLUM CNIDARIA

Nematocyts

- Tiny capsules made of chitin-like material and containing a coiled filament
- Filament may have tiny barbs or spines
- Some secrete an adhesive
- Touch causes nematocyst to discharge
- Nematocyst animation



© 1999 Addison Wesley Longman, Inc.

PHYLUM CNIDARIA - CNIDOCYTES

- *Mechanism of Nematocyst Discharge*
  - When stimulated, water rushes into the cnidocyte capsule
  - It opens, and nematocyst discharges
  - Barbs *inject poison* into prey

FIGURE 15.10



Animals from the phylum Cnidaria have stinging cells called cnidocytes. Cnidocytes contain large organelles called (a) nematocysts that store a coiled thread and barb. When hairlike projections on the cell surface are touched, (b) the thread, barb, and a toxin are fired from the organelle.

A CNIDOCYTE OF A HYDRA



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.



Table 33.1 Main Classes and Characteristics of the Cnidaria

Class and examples (est. # of species)	Class characteristics
Hydrozoa: *Obelia*, Portuguese man-of-war, *Hydra*, some corals (2,700)	Mostly marine; polyp stage usually dominant and colonial, reduced medusa stage
Scyphozoa: jellyfish (200)	All marine; medusa stage dominant and large (up to 2 m); reduced polyp stage
Anthozoa: sea anemones, sea fans, most corals (6,000)	All marine; polyp stage dominant; medusa stage absent; many are colonial
Cubozoa: box jellies, sea wasps (20)	All marine; medusa stage dominant; box-shaped

FIGURE 28.12 CLASS HYDROZOA



(a) *Obelia*

(b) *Physalia physalis* (Portuguese Man O' War)



(c) *Velella bae*

(d) *Hydra*

(a) Obelia, (b) Physalia physalis, known as the Portuguese Man O‘ War, (c) Velella bae, and (d) Hydra have different body shapes but all belong to the family Hydrozoa. (credit b: modification of work by NOAA; scale-bar data from Matt Russell)

Class Hydrozoa

Most are marine and colonial with **both** polyp and medusa forms

- *Hydra* is not typical (No medusa form)

Colonial *Obelia*

- Typical hydroid has a base, a stalk, and one or more individual polyp animals on “branches” which form feeding and reproductive structures.

The life cycle of the hydrozoan *Obelia*



HYDROZOAN DIVERSITY



Porpita



Velella



Obelia

CLASS HYDROZOAN

Hydra

- Found on the underside of aquatic leaves and lily pads in fresh water
- Worldwide distribution (16 species in North America)
- Body is a cylindrical tube
- Aboral end has a basal or pedal disc for attachment
- Ring of 6–10 hollow tentacles encircles mouth



CLASS HYDROZOAN

- The mouth opens to a gastrovascular cavity
- Buds may project from the side, each develop a mouth and tentacles
- Hydras feed on a variety of small crustaceans, insect larvae, and worms
  - Food organisms brush against the tentacles are captured by nematocysts
- Separate sexes (dioecious)
- Asexual budding



Hydra feeding



Small crustacean = Daphnia

© Cabisco/Visuals Unlimited



***Physalia* –**

Portuguese Man-o-war

MAN-O-WAR

PHYLUM CNIDARIA

Class Scyphozoa

Most of the larger jellyfishes belong to this class

Nearly all float in open sea

Bells vary in shape and size

- Composed mostly of mesoglea

Mesoglea contains amoeboid cells and fibers

Figure 9.16 *Aurelia* life history.



Scyphozoan life cycle



strobilation

STROBILATION

SCYPHOZOAN DIVERSITY


Larry S. Roberts

Aurelia **(moon jelly)**

Cyanea

Sexes are *separate*
- Gonads found in the gastric pouch of the female
- Zygote develops into a ciliated *planula larva*

PURPLE STRIPED JELLY, *PELAGIA PANOPYRA*



Lion's Mane jellyfish – 2 meters across and 20 meter tentacles



© R. Harbo

LION’S MANE JELLYFISH

PHYLUM CNIDARIA

Class Cubozoa



- Umbrella is square
  - One or more tentacles extend from each corner
- Strong swimmers
- Feed mostly on fish in nearshore areas
- The sea wasp (box jelly)
  - Potentially fatal sea wasp from Australia, heart failure and death in minutes. Most potent venom of all sea creatures.

FIGURE 15.14



A box jelly is an example from class Cubozoa.

FIGURE 28.11



(a) (b) (c)

The (a) tiny cubazoan jelly *Malo kingi* is thimble shaped and, like all cubozoan jellies, (b) has four muscular pedalia to which the tentacles attach. *M. kingi* is one of two species of jellies known to cause Irukandji syndrome, a condition characterized by excruciating muscle pain, vomiting, increased heart rate, and psychological symptoms. Two people in Australia, where Irukandji jellies are most commonly found, are believed to have died from Irukandji stings. (c) A sign on a beach in northern Australia warns swimmers of the danger. (credit c: modification of work by Peter Shanks)

DANGEROUS JELLYFISH

Only a *few jellyfish* (box jellyfish) *and the Portuguese man-of-war* can seriously harm humans.

Box jellyfish (“sea wasps”) have most powerful venom in the animal kingdom, affecting heart, nervous system and skin. Death by drowning of shock or heart attack is possible.

Found in Northern Australia in coastal waters. Most advanced jellyfish with clusters of 6 eyes on each side of the bell, with retina, iris and cornea (but no central nervous system!)

© Peter Parks/OSF/Animals Animals/Earth Scenes

Box jelly



OCELLI OF BOX JELLY



PHYLUM CNIDARIA

Class Anthozoa

Sea anemones and corals

No medusa stage

All marine, in both deep and shallow water, and vary in size

Fighting Anemones

ANTHOZOA – “flower animals” – no medusa stage







SEA ANEMONES

CORAL POLYPS



CLASS ANTHOZOA

Sea Anemones

- Polyps larger and heavier than hydrozoan polyps
- Attach to shells, rocks, timber, etc. by pedal discs
- *Venus flytrap anemone*

VENUS FLYTRAP ANEMONE



Figure 9.19 Class Anthozoa.



CLASS ANTHOZOA

- *When in danger,* water rapidly expelled through pores as the anemone contracts to a small size

- Most anemones can glide slowly on pedal discs
  - Some can swim with limited ability

Sea anemone escape from starfish



© R. Harbo

CLASS ANTHOZOA

- Most harbor symbiotic algae
- Some have a mutualistic relationship with hermit crabs or clown fish

Larry S. Roberts

CLASS ANTHOZOA

Reproduction

- Sexual – Some *dioecious* (male and female separate), some *monoecious* (hermaphrodites)
- Asexual - Small pieces of pedal disc break off and regenerate a small anemone

Corals



A B C

Larry S. Roberts

One coral polyp




Septal filaments
Mouth
Gastrovascular cavity
Gonad
Gastrodermal tube (solenia)
Coenenchyme
Tentacle
Axial rod
Pinnules

MYXOZOAN PARASITIC JELLYFISH RELATIVE IN YELLOWSTONE

Parasitic jellyfish? A jellyfish ancestor that has reduced itself to a parasitic lifestyle with a body of only a small number of cells.

Microscopic, they infect fish like the mountain whitefish and trout, causing kidney disease.

Some attack the spinal cartilage and nerves. Whirling disease makes fish swim in swirling corkscrew, or in circles chasing their tails. Can’t eat like this or get away from predators, so they starve or get eaten.




Cnidaria
Medusozoa
Anthozoa
Staurozoa
Scyphozoa
Cubozoa
Hydrozoa
Trachyline-like hydrozoa
Hydroids
Man-o-war
Other hydrozoa
Gut with septal filaments
Siphonoglyph
Anthozoan pharynx
Hexaradial and octaradial symmetry
Creeping planula without cilia
Strobilation
Complex eyes
Velarium
Boxlike medusa body
Rhopalium
Polyp reduced or lost
Polyp lost
Velum in medusae
Medusae produced by lateral budding and enntocodon
Medusoid body form
Motor nerve net
Primary polyp tentacles hollow
Mouth surrounded by solid tentacles
Planula larva
Cnidocytes
Radial, polypoid body form

PHYLUM CTENOPHORA ~ 100 species

- **CHARACTERISTICS – Similar to cnidarians, EXCEPT**
- **they have no nematocysts,**
- **have 8 rows of comb-like plates of modified cilia for locomotion, and**
- **have no sessile polyp stages**



Beroe

Pleurobrachia

A CTENOPHORE, OR COMB JELLY



SEA GOOSEBERRY, *PLEUROBRACHIA PILEUS*



CTENOPHORE



PHYLUM CTENOPHORA

Comb jellies

Less than 100 species – all marine and look like jellyfish

*First **complete gut** – mouth and anus (ctenophore eating)*

Eight rows of **cilia** on surface beat for propulsion

- Largest animals to use cilia for locomotion

Two long tentacles without stinging cells

Colloblasts secrete sticky substance

Hermaphroditic

Bioluminescent

Baroe eat other comb
jellies like Mnemiopsis

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Matthew J. D'Avella/SeaPics.com

PORIFERA – THE SPONGES

REVIEW OF ANIMAL PHYLOGENY


Porifera

Cnidaria and Ctenophora

Lophotrochozoa

Ecdysozoa

Protostomia

Deuterostomia

Radiata

Bilateria

Parazoa

Eumetazoa

Ancestral colonial choanoflagellate

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

FIGURE 15.8



© William C. Ober

Sponges are members of the phylum Porifera, which contains the simplest animals. (credit: Andrew Turner)


Red boring sponge
Encrusting sponge
Finger sponge
Variable sponge
Tube sponge
Coral head

PHYLUM PORIFERA

General Features

1. **Sessile** (“attached”) sponges are filter feeders

2. Porifera means “**pore-bearing**”

- Sac-like bodies perforated by many pores

3. Use flagellated “**collar cells**”, or *choanocytes,* to move water

- Body is efficient aquatic **filter**
  - Large sponges filter 1500 liters of water per day

Approximately 15,000 species of sponges

- *Most* are marine
  - A few live in brackish water, 150 in fresh water

PHYLUM PORIFERA - SHAPE OF LIFE VIDEO

1. Asymmetrical or superficially radially symmetrical
2. Three major cell types: pinacocytes (epidermal cells), amoeboid cells, and choanocytes
3. Central cavity (spongocoel), or a series of branching chambers, through which water circulates during filter feeding (Large sponges filter 1500 liters of water per day)
4. Have collagen. (Most abundant protein in all animals : 25%-35% of whole-body protein content.)
5. No tissues or organs

SHAPE OF LIFE SPONGES

SPONGES=PHYLUM PORIFERA

Despite their lack of complexity, sponges are clearly successful organisms, having persisted on Earth for more than **half a billion years**.

Sponge cells are arranged in a gelatinous matrix. Absence of true tissues or organs requires that all fundamental processes occur **at the level of individual cells**

Lacking a true digestive system, sponges depend on the **intracellular digestive processes** of their **choanocytes and amoebocytes** for their energy intake. The limit of this type of digestion is that food particles must be smaller than individual cells.

Gas exchange, circulation, and excretion occur by **diffusion** between cells and the water.

PHYLUM PORIFERA

Types of Canal Systems

Asconoid Sponges

- 1. *Simplest* body form (Asconoid)
- Small and tube-shaped
    - Center of the sponge (**spongocoel**) is covered with collar cell flagella
    - Water enters a large cavity, the spongocoel
        - Lined with *choanocytes*
        - Choanocyte flagella **pull water through** small holes (ostia) in cells called porocytes.

Morphology of a simple sponge.

Copyright © McGraw-Hill Education. Permission required for reproduction or display.



Choanocyte = collar cell



Exposed end has one *flagellum* surrounded by a *collar*

Collar

- Forms a *fine filtering device* to strain food
- Particles are trapped in mucous
- Food is moved to the *choanocyte* cell body and *engulfed (phagocytosis)*
- Food engulfed by choanocytes is passed to *amoeboid cells* for digestion

Copyright © 2016 McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education

PHYLUM PORIFERA

- *Amoeboid cells "Archaeocytes" Move about* in the mesohyl jelly matrix
  - Phagocytize particles
  - Can differentiate into any other type of cell,
  - Like a stem cell




© 1995 Jonathan Bird

Stem cell-like Amoebocytes

Epidermal

Amoebocyte – collagen producing



Gelatinous Ⓟ matrix = mesohyl

With collagen, other proteins, forming extracellular matrix

Secreted by sclerocytes (amoebocytes)

Sponge spicules





©DIAGRAM

A Ascon-type sponge (Class Calcarea)

B Skeleton of Venus's flower basket (Class Hexactinellida)

C Bath sponge (Class Demospongia)

D-F Calcareous spicules

G-J Siliceous spicules


BIODIDAC ©, J. Houseman

Calcareous spicules

Spongin



3 Sponge Classes – based on spicule composition

Hexactinellida Longest lived organisms on earth at 15,000 years per individual


BIODIDAC ©, J. Houseman

Calcarea



Demospongiae




Porifera
Calcarea
Demospongiae
Hexactinellida
Spicules not with 6 rays
Spongin network often present
Spicules with 6 rays
Calcium carbonate spicules
Siliceous spicules
Internal system of pores and canals for water flow

Choanoflagellate colony ancestor



Asconoid (*Leucosolenia*)



Syconoid (*Sycon*)

Ascon

Sycon



Leuconoid (*Euspongia*)

Complex sponges have increased surface area for choanocytes which results in filtering larger volumes of water = more food.

Leucon

PHYLUM PORIFERA

Most complex and larger sponges are called *Leucon* sponges

- Most sponges are leuconoid
- Have many oscula (holes)

- Much increased surface area with flagella compared to their volume
  - More collar cells can meet food demands
  - Large sponges filter 1500 liters of water per day

PHYLUM PORIFERA

Class Demospongiae

Contains *95% of living sponge species*

Spicules (if present) are made of silica

- Bound together by *spongin (protein network)*

All are *Leuconoid* body form

Bath sponges

- Lacks silica spicules
- Have spongin skeletons

PHYLUM PORIFERA

Sexual Reproduction

- Most sponges are *monoecious (both sexes in same individual)*
- Sperm sometimes arise from transformed *choanocytes, after meiosis*
- Eggs arise from amoebocytes

PHYLUM PORIFERA

- In some, one sponge releases sperm which enter the pores of another sponge
- Some sponges release both sperm and oocytes into water
- Form a *free-swimming larva*

Sexual and asexual reproduction

Embryos are free-swimming, *adult sponges* always attached



SPONGE LARVAE





The azure vase-sponge *Callyspongia plicifera* is a type of sponge that releases short-lived (minutes only) larvae that settle close to the parent 0.25X

PHYLUM PORIFERA

- Sponges have a great ability to regenerate lost parts and repair injuries
- Put them in a blender and they can reassemble themselves
- Regeneration after fragmentation is one means of asexual reproduction

PHYLUM PORIFERA

- Asexual reproduction can also occur by *bud formation*
    - *External buds*
        - Small individuals that break off after attaining a certain size
    - *Gemmules*
        - Formed by amoebocytes (archaeocytes) coated with tough spongin and spicules
        - Survive harsh environmental conditions

GEMMULES – sponge asexual reproduction



Gemmule


Micropyle
Inner membrane
Archaeocytes
Spicules


spicules
gemmule
© W.P. Armstrong 2003

ECOLOGY

Many live as *commensals (or parasites)* in or on other sponges

- Also grow on a variety of other living organisms

Have few predators

- Sponges may have an *elaborate skeletal structure* and often have a *noxious odor*
- Sea turtles eat them.

FAMILY CLADORHIZIDAE

(DEMOSPONGIAE)

Carnivorous sponge that preys on crustaceans and other small animals, rather than filter feeding! Perhaps an adaptation to living in very deep waters with low nutrient levels (oligotrophic).





CARNIVOROUS SPONGES

CLADORHIZIDAE 1



JONATHAN BIRD
SPONGE VIDEO 9 MIN

CHAPTER 32

AN INTRODUCTION TO ANIMAL DIVERSITY

Key Concepts:

Characteristics of Animals

Animal Classification

Evolutionary History of Animal Diversity

Characteristics of Animals

Multicellular **heterotrophs**

Cells **lack cell walls**

Most have nerves, muscles, and capacity to **move** at some point in the life cycle

Able to **reproduce sexually**

Specialized sensory structures and **nervous system**

Cells exist in extensive **extracellular matrix**

Unique **cell junctions**

All animals have **Hox genes** and small ribosomal subunit **rRNA**

FIGURE 27.5



Hox genes are highly conserved genes encoding transcription factors that determine the course of embryonic development in animals. In vertebrates, the genes have been duplicated into four clusters: *Hox-A*, *Hox-B*, *Hox-C*, and *Hox-D*. Genes within these clusters are expressed in certain body segments at certain stages of development. Shown here is the homology between *Hox* genes in mice and humans. Note how Hox gene expression, as indicated with orange, pink, blue and green shading, occurs in the same body segments in both the mouse and the human.

Animal Classification

Most biologists agree kingdom is **monophyletic**

About 35 recognized animal phyla

Most likely ancestor was a colonial flagellated protist similar to **choanoflagellates**

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Kingdom Animalia

Metazoa

Parazoa | Eumetazoa

Radiata | Bilateria

Protostomia | Deuterostomia

Lophotrochozoa | Ecdysozoa

Porifera

Cnidaria and Ctenophora

Platyhelminthes

Rotifera

Bryozoa

Brachiopoda

Mollusca

Annelida

Nematoda

Arthropoda

Echinodermata

Chordata

Most with lophophore or trochophore larva

Ecdysis

Protostome development

Deuterostome development, endoskeleton

Bilateral symmetry

Radiata

Parazoa

Tissues

Multicellularity

Common ancestor of animals and choanoflagellates

KEY

■ Critical innovations



Metazoa (animals)

Eumetazoa (specialized tissues)

Bilateria (bilateral symmetry, triploblastic)

Possess body cavity

Protostomia

Ecdysozoa

Lophotrochozoa

Deuterostomia

Acoelomates (no coelom)

Radiata (radial symmetry, diploblastic)

Parazoa (no tissues)

Arthropoda (arthropods)

Nematoda (roundworms)

Platyhelminthes (flatworms)

Rotifera (rotifers)

Ectoprocta

Brachiopoda

Annelida (annelids)

Mollusca (mollusks)

Chordata (chordates)

Echinodermata (echinoderms)

Acoela

Cnidaria (cnidarians)

Ctenophora (comb jellies)

Porifera (sponges)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Choanoflagellate cell

Sponge cell (choanocyte)

(a) Colonial choanoflagellate

(b) Sponge

TRADITIONAL CLASSIFICATION BASED ON BODY PLANS

Morphological and developmental features traditionally used to classify animals:

1. Presence or absence of different **tissue types**
2. Type of **body symmetry**
3. Specific features of **embryonic development**

1. **Tissues**

- **Metazoa** – all animals
  - Metazoa divided based on whether there are specialized tissues
- **Parazoa** (without specialized tissues or organs)
  - **Porifera** – sponges (“beside” animals)
- **Eumetazoa** (more than one type of tissue and organ)
  - **All other animals**

2. Symmetry

- **Eumetazoa** are radially symmetrical or bilaterally symmetrical
  - **Bilateria** (bilateral animals)
    - Have cephalization and **dorsal** and **ventral** sides
    - 3 germ layers
  - **Radiata** (radial animals)
    - Have **oral** and **aboral** sides
    - 2 germ layers

FIGURE 27.7



The (a) sponge (Parazoa) is asymmetrical. The (b) jellyfish and (c) anemone are radially symmetrical (Eumetazoa, 2 tissue layers), and the (d) butterfly is bilaterally symmetrical (Eumetazoa, 3 tissue layers). (credit a: modification of work by Andrew Turner; credit b: modification of work by Robert Freiburger; credit c: modification of work by Samuel Chow; credit d: modification of work by Cory Zanker)

FIGURE 27.8



The bilaterally symmetrical human body can be divided into planes.

(a) Parazoa: no tissue types





(b) Eumetazoa: two tissue types
Radiata: radial symmetry

(c) Eumetazoa: three tissue types
Bilateria: bilateral symmetry



Number of cell layers

- **Bilateria are triploblastic** – 3 layers
- **Radiata are diploblastic** – 2 layers
- Cell layers develop during gastrulation
- Inner layer – endoderm
- Outer layer – ectoderm
- Mesoderm - 3rd layer in bilateral animals
  - Forms muscles and most other organs

FIGURE 27.9



During embryogenesis, diploblasts develop two embryonic germ layers: an ectoderm and an endoderm. Triploblasts develop a third layer—the mesoderm—between the endoderm and ectoderm.

FIGURE 27.4



During embryonic development, the zygote undergoes a series of mitotic cell divisions, or cleavages, to form an eight-cell stage, then a hollow blastula. During a process called gastrulation, the blastula folds inward to form a cavity in the gastrula.



FIGURE 27.10



(a) **Acoelomate** (flatworms)

(b) **Eucoelomate** (annelids, mollusks, arthropods, echinoderms, chordates)

(c) **Pseudocoelomate** (roundworms)

Triploblasts may be (a) acoelomates, (b) eucoelomates, or (c) pseudocoelomates. Acoelomates have no body cavity. Eucoelomates have a body cavity within the mesoderm, called a coelom, which is lined with mesoderm. Pseudocoelomates also have a body cavity, but it is sandwiched between the endoderm and mesoderm. (credit a: modification of work by Jan Derk; credit b: modification of work by NOAA; credit c: modification of work by USDA, ARS)


Zygote
Blastula
Gastrula
Ectoderm (outer layer)
Mesoderm (middle layer)
Endoderm (internal layer)
Germ cells
Outer surface
Central nervous system
Neural crest
Dorsal
Paraxial
Intermediate
Lateral
Head
Digestive tube
Pharynx
Respiratory tube
Male
Female
Epidermal cells of skin
Neuron of brain
Pigment cell (melanocyte)
Notochord
Bone tissue
Tubule cell of the kidney
Red blood cells
Facial muscle
Stomach cell
Thyroid cell
Lung cell (alveolar cell)
Sperm
Egg

3. **Embryonic development in Eucoelomates**

- ❑ **Protostome**
  - Spiral cleavage
  - Cleavage is determinate
  - Blastopore becomes mouth
- ❑ **Deuterostome**
  - Radial cleavage
  - Cleavage is indeterminate – pluripotent stem cells
  - Blastopore becomes anus

FIGURE 27.11

Eucoelomates can be divided into two groups based on their early embryonic development. In protostomes, part of the mesoderm separates to form the coelom in a process called schizocoely. In deuterostomes, the mesoderm pinches off to form the coelom in a process called enterocoely. It was long believed that the blastopore developed into the mouth in protostomes and into the anus in deuterostomes, but recent evidence challenges this belief.



A COMPARISON OF EARLY DEVELOPMENT IN PROTOSTOMES AND DEUTEROSTOMES



Protostomes (mollusks, annelids, arthropods)
Deuterostomes (echinoderms, chordates)
(a) Cleavage
Eight-cell stage
Spiral and determinate
Eight-cell stage
Radial and indeterminate
(b) Coelom formation
Coelom
Archenteron
Coelom
Mesoderm
Blastopore
Blastopore
Mesoderm
Schizocoelous: solid masses of mesoderm split to form coelom
Enterocoelous: folds of archenteron form coelom
(c) Fate of blastopore
Anus
Mouth
Digestive tube
Mouth
Anus
Mouth develops from blastopore
Anus develops from blastopore

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

OTHER METHODS OF CLASSIFICATION

Possession of **exoskeleton**

Development of **notochord**

Presence or absence of **segmentation**

- Traced to changes in homeotic or *Hox* genes

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



In earthworms, each ring is a distinct segment.

Lobsters have developed specialized appendages on many segments.

Fishes exhibit segmentation in their muscles and backbone.

SIMILARITIES BETWEEN TRADITIONAL AND MOLECULAR PHYLOGENY

1. The clade called **Metazoa** is monophyletic, meaning all animals came from a single common ancestor.
2. At the earliest stages of evolution, molecular phylogeny supports the traditional view of the split between **Parazoa** and **Eumetazoa.**
3. There is also agreement about an early split between **Radiata** and **Bilateria,** with most animal phyla belonging to the Bilateria.
4. Molecular phylogeny also agrees that the echinoderms and chordates belong to a clade called the **Deuterostomia.**

DIFFERENCES IN THE NEW MOLECULAR PHYLOGENY

Two new invertebrate clades:

Ecdysozoa

  - Named for ecdysis (molting)
  - **Nematodes, Arthropods,** and a few other minor phyla

Lophotrochozoa

  - Named for the lophophore (feeding tentacles) and trochophore larva
  - **Mollusks, Annelids,** and several other phyla

ECDYSIS



Benjamin Cummings

A LOPHOPHORATE

Lophophore



A TROCHOPHORE LARVA



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

© Dwight Kuhn

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) Lophophore of a phoronid worm

(b) Trochophore larva

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Table 32.2 Summary of the Basic Characteristics of the Major Animal Phyla

Feature	Porifera (sponges)	Cnidaria and Ctenophora (hydra, anemones, jellyfish)	Platyhelminthes (flatworms)	Rotifera (rotifers)	Lophophorata (bryozoans and others)	Mollusca (snails, clams, squids)	Annelida (segmented worms)	Nematoda (roundworms)	Arthropoda (insects, arachnids, crustaceans)	Echinodermata (sea stars, sea urchins)	Chordata (vertebrates and others)
Estimated number of species	8,000	11,000	20,000	2,000	4,000+	110,000	15,000	20,000	1,000,000+	7,000+	52,000+
Level of organization	Cellular; lack tissues and organs	Tissue; lack organs	Organs	Organs	Organs	Organs	Organs	Organs	Organs	Organs	Organs
Symmetry	Absent	Radial	Bilateral	Bilateral	Bilateral	Bilateral	Bilateral	Bilateral	Bilateral	Bilateral larvae, radial adults	Bilateral
Cephalization	Absent	Absent	Present	Present	Reduced	Present	Present	Present	Present	Absent	Present
Germ layers	Absent	Two	Three	Three	Three	Three	Three	Three	Three	Three	Three
Body cavity	Absent	Absent	Absent	Pseudocoelom	Coelom	Reduced coelom	Coelom	Pseudocoelom	Reduced coelom	Coelom	Coelom
Segmentation	Absent	Absent	Absent	Absent	Absent	Absent	Present	Absent	Present	Absent	Present (but reduced)

ANIMAL PHYLOGENY BASED ON SEQUENCING OF SSU-RRNA



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.



THE MAJOR GROUPS OF ANIMALS THROUGH TIME

The width of each oval suggests the relative number of species in that group

Hickman Zoology

DOMAIN EUKARYA: PROTISTS



Protists range from the microscopic, single-celled (a) *Acanthocystis turfacea* and the (b) ciliate *Tetrahymena thermophila*, both visualized here using light microscopy, to the enormous, multicellular (c) kelps (Chromalveolata) that extend for hundreds of feet in underwater “forests.” (credit a: modification of work by Yuiuji Tsukii; credit b: modification of work by Richard Robinson, Public Library of Science; credit c: modification of work by Kip Evans, NOAA; scale-bar data from Matt Russell)

CHARACTERISTICS OF EUKARYOTES

1. Nuclei with nuclear envelope with nuclear pores.
2. Mitochondria (ancestor of mitochondria an alphaproteobacteria)
3. Cytoskeleton w/ actin microfilaments and microtubules
4. Flagella and cilia, maybe pseudopodia
5. Chromosomes with protein histones
6. Mitosis
7. Sex with meiosis and haploid nuclei
8. Cell walls found in some organisms of all lineages of protozoa
9. Ancestor a composite of host cell (probably an archaea) and cell of alpha-proteobacterium inside (**endosymbiosis**)

PROTOZOA

The “protists”

Catch-all term for the non-plant, non-animal, non-fungal eukaryotes. No taxonomic or systematic significance.

More diverse than the rest of the eukaryotes. Like multiple kingdoms

(Unicellular, multicellular, multin
walls, cell membranes, pellicle)

INTRODUCTION TO PROTISTS VIDEO

PROTISTS

Protist video 4:11

Key Topics:

- **An Introduction to Protists**
- **Nutritional and Defensive Adaptations**
- **Reproductive Adaptations**
- **Evolution and Relationships**

OLD PROTIST VIDEO: (NO LONGER CALLED “KINGDOM PROTISTA”)



An Introduction to Protists

Protists are eukaryotes that are not classified in the plant, animal, or fungal kingdoms, though some protists are closely related to plants or animals or fungi.

Two common characteristics

- ❑Most abundant in moist habitats
- ❑Most of them are microscopic in size

CLASSIFIED BY ECOLOGICAL ROLE

Three major groups:

- **Algae** – generally photoautotrophic
- **Protozoa** – heterotrophic
- **Fungus-like** – resemble fungi in body form and absorptive nutrition

Terms lack taxonomic or evolutionary meaning

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Photographs by H. Cantor-Lund reproduced with permission of the copyright holder J. W. G. Lund

ALGAL PROTISTS

Variety of **pigments**

- ☐ Adapt photosystems to capture more light
- ☐ Water absorbs the longer red and yellow wavelengths more than the shorter blue and green wavelengths
- ☐ Accessory pigments absorb light and transfer it to chlorophyll a

Variety of types of **food storage molecules**

- ☐ Starch, polysacchrides, and oil

NUTRITION

Protists, the most nutritionally diverse of all eukaryotes, include:

- **Autotrophs**, which contain chloroplasts



- **Heterotrophs**, which absorb organic molecules or ingest larger food particles through phagocytosis – an infolding of the plasma membrane around a food particle

- **Mixotrophs**, which combine photosynthesis and heterotrophic nutrition



phagocytosis

Dinobryon – a golden alga

Photosynthetic stramenopile that lives in the phytoplankton of freshwater lakes

Example of a **mixotroph**

Switch back and forth between photoautotrophy and heterotrophy, depending on conditions in their environment

Remarkable **nutritional flexibility**



CC BY-SA 3.0 de

FIGURE 13.14 PHAGOCYTOSIS


Phagocytosis
Food particle
Food vacuole
Exocytic vesicle containing undigested material
Phagocytosis
Pseudopods
Lysosome containing digestive enzymes

The stages of phagocytosis include the engulfment of a food particle, the digestion of the particle using hydrolytic enzymes contained within a lysosome, and the expulsion of undigested material from the cell.

CYTOPHARYNX

Some protozoa ingest food in a specialized region called a cytopharynx.



CLASSIFIED BY HABITAT

Particularly common and diverse in oceans, lakes, wetlands and rivers

Plankton – swimming or floating

- **Phytoplankton** – photosynthetic
- **Zooplankton** – heterotrophic
- Occur primarily as single cells, colonies or short filaments

Also found in:

Soil
Snow
Dead organisms or waste (decomposers)
As parasites inside other organisms

Phytoplankton



CLASSIFIED BY MOTILITY

Swim using eukaryotic flagella

- **Flagellates**
- Some flagellated reproductive cells

Cilia – shorter and more abundant than flagella, and beat in synchrony

- **Ciliates**

Amoeboid movement – using pseudopodia

- **Amoebae**

Gliding on protein or carbohydrate slime

FORM AND FUNCTION CONTINUED

Locomotor Organelles

Cilia and **Flagella** – both have similar internal structure. 9 pairs of longitudinal microtubules arranged in a circle around a central pair.

9+2 arrangement of microtubules.

Cilia Video

More cilia



CILIA



CILIATED PROTISTS



Visualization of a 3D reconstructed sea urchin sperm flagellum.



Flagella Video

Nicastro D et al. PNAS 2005;102:15889-15894

©2005 by National Academy of Sciences

PNAS

Flagellated protists

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Dr. Dennis Microscopy/Visuals Unlimited

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Dr. Stanley Flegler/Visuals Unlimited

PSEUDOPODIA

Protrusions of the cytoplasm, usually in amoeba, that are used for both locomotion and feeding. Video

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



CYTOPLASMIC STREAMING



OTHER TYPES OF PSEUDOPODS

Long, thin pseudopodia with multiple rods of microtubules, as in Radiolarians, Foraminiferans

Actinopods.



CC0

ACTINOPODS



Reproductive Adaptations

Asexual reproduction

- All protists can reproduce asexually
- Many produce cysts with thick, protective walls that remain dormant in bad conditions
- Many protozoan pathogens spread from one host to another via cysts

ASEXUAL REPRODUCTION

Asexual

- **binary fission**, splitting into 2 cells, similar to mitosis, both cells similar size.
- **budding** – much smaller cell breaks off larger one.

All protists can reproduce asexually.

Volvox produces daughter colonies asexually

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





EVOLUTION OF VOLVOX 0.11, 0.45

VOLVOX COLONIES WITH DAUGHTER COLONIES



SEXUAL REPRODUCTION

Eukaryotic sexual reproduction with gametes and zygotes arose among the protists

Generally adaptive because it produces diverse genotypes

Sporic life cycle

- Many multicellular green and brown seaweeds
- Also known as **alternation of generations**
- Two types of multicellular organisms
  - **Haploid gametophyte produces gametes by mitosis**
  - **Diploid sporophyte produces spores by meiosis**

THE LIFE CYCLE OF *LAMINARIA* (BROWN ALGA): AN EXAMPLE OF ALTERNATION OF GENERATIONS



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Ciliate sexual reproduction

- Most complex sexual process in protists
- Have two types of nuclei (single **macronucleus** and one or more **micronuclei**)
- Macronuclei are the source of the information for cell function
- Two cells pair and fuse – **conjugation**

Sexual reproduction involves a meiosis event of the DNA in the **micronucleus**.

This haploid DNA is exchanged with another organism, restoring a diploid condition.



REPRODUCTION

Sexual reproduction

Provides new combinations of genetic material

But protozoa do not have “embryos”

Haploid DNA (gamete nuclei) is exchanged between individuals

It fuses with the gamete nuclei of the other organism to become diploid, (may go through meiosis soon to return to haploid condition). Many protists are haploid.



PARAMECIUM CONJUGATING



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

23.16



1 Two different mating types form a cytoplasmic bridge.

2 Meiosis produces four haploid micronuclei, three of which disintegrate.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)



1 Two different mating types form a cytoplasmic bridge.

2 Meiosis produces four haploid micronuclei, three of which disintegrate.

3 Each remaining micronucleus divides by mitosis.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)


Macronucleus
Cytoplasmic bridge
Micronucleus
2
Meiosis produces four haploid micronuclei, three of which disintegrate.
cium
3
Each remaining micronucleus divides by mitosis.
4
The conjugate pair swaps micronuclei.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)



1 Two different mating types form a cytoplasmic bridge.

2 Meiosis produces four haploid micronuclei, three of which disintegrate.

3 Each remaining micronucleus divides by mitosis.

4 The conjugate pair swaps micronuclei.

5 The haploid micronuclei fuse, forming a diploid micronucleus.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)



Reproduction in *Paramecium*

1 Two different mating types form a cytoplasmic bridge.

2 Meiosis produces four haploid micronuclei, three of which disintegrate.

3 Each remaining micronucleus divides by mitosis.

4 The conjugate pair swaps micronuclei.

5 The haploid micronuclei fuse, forming a diploid micronucleus.

6 The micronucleus undergoes three rounds of mitosis, producing eight micronuclei.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)



Sexual Reproduction in *Paramecium*

1 Two different mating types form a cytoplasmic bridge.

2 Meiosis produces four haploid micronuclei, three of which disintegrate.

3 Each remaining micronucleus divides by mitosis.

4 The conjugate pair swaps micronuclei.

5 The haploid micronuclei fuse, forming a diploid micronucleus.

6 The micronucleus undergoes three rounds of mitosis, producing eight micronuclei.

7 The original macronucleus disintegrates, and four of the micronuclei become macronuclei.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)



Sexual Reproduction in *Paramecium*

1 Two different mating types form a cytoplasmic bridge.

2 Meiosis produces four haploid micronuclei, three of which disintegrate.

3 Each remaining micronucleus divides by mitosis.

4 The conjugate pair swaps micronuclei.

5 The haploid micronuclei fuse, forming a diploid micronucleus.

6 The micronucleus undergoes three rounds of mitosis, producing eight micronuclei.

7 The original macronucleus disintegrates, and four of the micronuclei become macronuclei.

8 Two rounds of cell division produce four daughter cells.

The complex process of sexual reproduction in *Paramecium* creates eight daughter cells from two original cells. Each cell has a macronucleus and a micronucleus. During sexual reproduction, the macronucleus dissolves and is replaced by a micronucleus. (credit “micrograph”: modification of work by Ian Sutton; scale-bar data from Matt Russell)

Evolution and Relationships

At one time, protists were in a single kingdom (not since 1990)

- However, “protists” is **not a monophyletic group**

Evolutionary understanding is **in flux**

- Some relationships are uncertain or disputed
- New protists still being discovered

Classified into **supergroups**



This diagram shows a proposed classification of the domain Eukara. Currently, the domain Eukarya is divided into six supergroups. Within each supergroup are multiple kingdoms. Dotted lines indicate suggested evolutionary relationships that remain under debate.

SUPERGROUP EXCAVATA

Related to some of Earth’s earliest eukaryotes

Named for **a feeding groove** “excavated” into the cells of many representatives

Food particles are taken into cells by **phagotrophy**

  - Endocytosis
  - Evolutionary basis for endosymbiosis

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Bacterial prey are consumed by the process of phagocytosis.

1 Prey is enveloped by the plasma membrane at the feeding groove.

2 The membrane pinches off a food vesicle within the cytoplasm.

3 Other vesicles deliver digestive enzymes to food vesicles; prey is digested for food.

(a) Excavate *Jakoba* with feeding groove

(b) The process of phagocytosis in excavates

GIARDIA INTESTINALIS, A DIPLOMONAD



Excavata

Some are parasites

Trichomonas vaginalis, Giardia intestinalis

Once thought to lack mitochondria

Possess highly modified mitochondria

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) *Trichomonas vaginalis*



(b) *Giardia intestinalis*

a: © Dr. David M. Phillips/Visuals Unlimited; b: © Jerome Paulin/Visuals Unlimited

TRICHOMONAS VAGINALIS, A PARABASALID (EXCAVATA)



HYPERMASTIGID

Another parabasalid (like *Trichomonas*) but not parasitic. It is found in the gut of termites and cockroaches, necessary to break down wood that is consumed.

SUPERGROUP EXCAVATA

Euglenozoa

- Protein strips under plasma membrane allow crawling
- Some are heterotrophic, but *Euglena* is photosynthetic

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) *Euglena*

a: © Wim van Egmond/Visuals Unlimited

EUGLENA MOVEMENT

***TRYPANOSOMA BRUCEI*, THE EUGLENOZOAN (KINETOPLASTID) THAT CAUSES SLEEPING SICKNESS**



Benjamin Cummings

SUPERGROUP EXCAVATA: EUGLENOZOA

Kinetoplastids

- Named for unusually large mass of DNA (kinetoplast) in a single large mitochondrion
- *Leishmania*
- *Trypansosoma brucei*

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(b) ***Leishmania*** **(c)** ***Trypanosoma***

b: © Ross Waller, University of Melbourne, Australia;
c: © Eye of Science/Photo Researchers, Inc.



This diagram shows a proposed classification of the domain Eukara. Currently, the domain Eukarya is divided into six supergroups. Within each supergroup are multiple kingdoms. Dotted lines indicate suggested evolutionary relationships that remain under debate.

SUPERGROUP ALVEOLATA (“SAR”)

Named for **saclike membranous vesicles** (alveoli) present in cell periphery

Ciliophora

- Ciliates – conjugation

Apicomplexa

- ALL are intracellular parasites e.g. P*lasmodium* (causative agent of malaria)

Dinoflagellates

- Some photosynthetic, others not
- Red tide and mutualistic relationship with coral

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Alveolata

Ciliophora

Apicomplexa

Dinozoa

Stramenopila

Oomycota

Bacillariophyceae

Phaeophyceae

Rhizaria

Chlorarachniophyta

Radiolaria

Foraminifera

Alveoli

Flagellar hairs

Filose pseudopodia

KEY

- 🟩 Algae
- 🟧 Protozoa
- 🟦 Fungus-like protists
- ⬛ Critical innovations

CHROMOALVEOLATA: CILIATES: *STENTOR* (LEFT), *PARAMECIUM* (RIGHT)



CILIATES: *PARAMECIUM*



(c) *Paramecium*

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Contractile vacuole

SPECIALIZED ORGANELLES - CILIATES

Contractile Vacuoles Responsible for water balance or **osmoregulation**

Found mainly in freshwater protozoa, whose cellular water concentration is lower than the environment

Act as a pump to remove excess water from the cytoplasm



CONTRACTILE VACUOLES



PARASITIC PROTIST LIFE CYCLE - APICOMPLEXA

Parasitic protists often use **more than one host organism,** in which different life stages occur

ex: **Malarial parasite *Plasmodium***

Alternates between humans and *Anopheles* mosquitoes

Different stages in different hosts and host tissues

Chromoalveolata: Apicomplexan: *Plasmodium* in red blood cells



Wellcome Images

PLASMODIUM LIFE CYCLE

In human

In mosquito

Plasmodium (Malaria) Life Cycle in humans



Plasmodium (Malaria) life cycle in Mosquitos



THE TWO-HOST LIFE HISTORY OF *PLASMODIUM*, THE APICOMPLEXAN THAT CAUSES MALARIA



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.



This diagram shows a proposed classification of the domain Eukara. Currently, the domain Eukarya is divided into six supergroups. Within each supergroup are multiple kingdoms. Dotted lines indicate suggested evolutionary relationships that remain under debate.

HYPERMASTIGID

Another parabasalid (like *Trichomonas*) but not parasitic. It is found in the gut of termites and cockroaches, necessary to break down wood that is consumed.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Alveolata

Ciliophora

Apicomplexa

Dinozoa

Stramenopila

Oomycota

Bacillariophyceae

Phaeophyceae

Rhizaria

Chlorarachniophyta

Radiolaria

Foraminifera

Alveoli

Flagellar hairs

Filose pseudopodia

KEY

- 🟩 Algae
- 🟧 Protozoa
- 🟦 Fungus-like protists
- ⬛ Critical innovations


i = Infective Stage
d = Diagnostic Stage
CDC
SAFER·HEALTHIER·PEOPLE™
Human Liver Stages
Liver cell
Infected liver cell
②
③
Schizont
④ Ruptured schizont
A
Exo-erythrocytic Cycle
Mosquito Stages
⑫ Ruptured oocyst
Release of sporozoites
⑪ Oocyst
① Mosquito takes a blood meal (injects sporozoites)
C
Sporogonic Cycle
⑩ Ookinete
⑧ Mosquito takes a blood meal (ingests gametocytes)
Macrogametocyte
Microgamete entering macrogamete ⑨
Exflagellated microgametocyte
Human Blood Stages
⑤
Immature trophozoite (ring stage)
B
Erythrocytic Cycle
Mature trophozoite
⑥ Ruptured schizont
Schizont
⑦ Gametocytes
P. falciparum
♀
♂
⑦ Gametocytes
♀
♂
P. vivax
P. ovale
P. malariae

ALVEOLATA: DINOFLAGELLATE



SWIMMING WITH BIOLUMINESCENT DINOFLAGELLATES



Benjamin Cummings

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



SUPERGROUP STRAMENOPILA ("SAR")

Named for distinctive **strawlike hairs** on the surface of flagella

Wide range of algae, protozoa, and fungus-like protists

Usually produce **flagellated cells** at some point

Heterotrophic or photosynthetic

  - Plastids from secondary endosymbiosis with red algae

- Includes the Oomycetes (water molds), Brown algae, Golden algae and diatoms

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Flagellar hairs

Plastids

Nucleus

Food storage

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) Diatom



(b) Kelp forest

a: © Linda Graham; b: © Jeff Rotman/Photo Researchers, Inc.



DIATOM SHELL



KELP FOREST (*MACROCYSTIS*)



Benjamin Cummings

KELP FOREST (*MACROCYSTIS*)



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Alveolata

Ciliophora

Apicomplexa

Dinozoa

Stramenopila

Oomycota

Bacillariophyceae

Phaeophyceae

Rhizaria

Chlorarachniophyta

Radiolaria

Foraminifera

Alveoli

Flagellar hairs

Filose pseudopodia

KEY

🟩 **Algae**

🟧 **Protozoa**

🟦 **Fungus-like protists**

⬛ **Critical innovations**

SUPERGROUP RHIZARIA (“SAR”)

Have thin, hairlike extensions of the cytoplasm

Phylum Radiolaria

Phylum Foraminifera

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) Radiolarian



(b) Foraminiferan

a: © Claude Nuridsany & Marie Perennou/SPL/Photo Researchers, Inc.; b: © O. Roger Anderson, Columbia University,
Lamont-Doherty Earth Observatory

ARCHAEPLASTIDA: PLANTS AND RELATIVES

Supergroup that includes land plants also encompasses several algal phyla

Kingdom Plantae (land plants) evolved from green algal ancestors

Phylum Chlorophyta and Charophyta – green algae

Phylum Rhodophyta – red algae

GREEN ALGAE

Phylum Chlorophyta

Diverse structural types

Occur in fresh water, the ocean, and on land

Cells contain same type of plastids and photosynthetic pigments as in land plants.

Phylum Charophyta

Closest relatives to land plants

Similar reproductive patterns and morphology

RED ALGAE

Most are multicellular marine macroalgae

Red appearance due to **distinctive photosynthetic pigments**

Lack flagella

Unusually complex life cycles

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) ***Calliarthron***

(b) ***Chondrus crispus***

a: © Doug Sokell/Visuals Unlimited; b: © Andrew J. Martinez/Photo Researchers, Inc.

SUPERGROUP AMOEBOZOA

Many types of amoebae

Move using **pseudopodia**

ex: ***Dictyostelium discoideum,*** Slime Molds

- Model organism for understanding movement, cell communication, and development.
- In response to starvation, single amoebae aggregate into a multicellular “slug” that develops into a stalked structure containing spores
- Spores pop out and produce new amoebae

SLIME MOLD



DICTYOSTELIUM LIFE CYCLE



THE LIFE CYCLE OF A CELLULAR SLIME MOLD (*DICTYOSTELIUM*)



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

FIGURE 23.28


Cellular Slime Mold Life Cycle
Germination gives rise to amoeboid cells.
Spore (1n)
The spores disperse.
Mitosis and Meiosis
Asexual reproduction
Solitary amoebas (1n)
Zygote (2n)
Sexual reproduction
Fertilization
Mitosis without cytokinesis results in multiple solitary amoebas that may aggregate. The aggregated amoebas are able to migrate.
Aggregated amoebas
Fruiting body
Stalk
Cells migrate up the stalk and form spores.
A stalk forms with a fruiting body at the top.

Cellular slime molds may exist as solitary or aggregated amoebas. (credit: modification of work by “thatredhead4”/Flickr)

STAGES OF *DICTYOSTELIUM*



FIGURE 23.27


Plasmodial Slime Mold Life Cycle
Germination gives rise to cells that can convert between amoeboid and flagelated forms.
Spores (1*n*)
Meiosis within the sporangium results in haploid spores.
Sporangium
Stalk
Amoeboid
Flagellated
Fertilization by fusion of either cell type results in a 2*n* zygote.
The plasmodium matures and sporangia form.
Mature plasmodium
Zygote (2*n*)
Mitosis without cytokinesis results in a single- celled multinucleate mass called a plasmodium
Feeding plasmodium

The life cycle of the plasmodial slime mold is shown. The brightly colored plasmodium in the inset photo is a single-celled, multinucleate mass. (credit: modification of work by Dr. Jonatha Gott and the Center for RNA Molecular Biology, Case Western Reserve University)

DOG VOMIT SLIME MOLD



PRETZEL SLIME MOLD



SUPERGROUP OPISTHOKONTA

Includes animal and fungal kingdoms and related protists

Named for single posterior flagellum on swimming cells

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Stephen Fairclough, King Lab, University of California at Berkeley

Choanoflagellate protists

- Feature distinctive collar surrounding flagella
- These are the modern protists most related to the common ancestor of animals

Figure 22.1



- Certain prokaryotes can live in extreme environments such as the Morning Glory pool, a hot spring in Yellowstone National Park. The spring’s vivid blue color is from the prokaryotes that thrive in its very hot waters. (credit: modification of work by Jon Sullivan)

Archaea and Bacteria - Prokaryotes

Key Topics:

- **Diversity and Evolution**
- **Structure and Movement**
- **Reproduction**
- **Nutrition and Metabolism**
- **Ecological Roles and Biotechnological Applications**

Diversity and Evolution

- One of the most prominent features of the bacteria and archaea is their diversity
- Domain **Archaea**
- Domain **Bacteria** (also called Eubacteria)
- Both prokaryotic – lack nucleus



Universal Ancestor

Domain Eukarya

Domain Archaea

- Euryarchaeotes
- Crenachaeotes
- Nanoarchaeotes
- Korarchaeotes

Domain Bacteria

- Proteobacteria
- Chlamydias
- Spirochetes
- Cyanobacteria
- Gram-Positive bacteria

½ the earth's biomass is prokaryotic!



Bacteria on the point of a pin



Figure 22.2



- This microbial mat, about one meter in diameter, grows over a hydrothermal vent in the Pacific Ocean in a region known as the “Pacific Ring of Fire.” The mat helps retain microbial nutrients. Chimneys such as the one indicated by the arrow allow gases to escape (credit a: modification of work by Dr. Bob Embley, NOAA PMEL, Chief Scientist; credit b: modification of work by Ricardo Murga, Rodney Donlan, CDC; scale-bar data from Matt Russell)

Figure 22.3



(a) These living stromatolites are located in Shark Bay, Australia.

(b) These fossilized stromatolites, found in Glacier National Park, Montana, are nearly 1.5 billion years old. (credit a: Robert Young; credit b: P. Carrara, NPS)

Domain Archaea

- Have features in common with the eukaryotic nucleus and cytoplasm, suggesting common ancestry
  - Histone proteins associated with DNA
  - 33 Ribosomal proteins shared
  - RNA polymerases closely related to those of eukaryotes
- Unique membrane lipids give resilience
  - **Ether**-bonded lipids more resistant to heat and other conditions
  - Allow Archaea to live in extreme environments
  - “Extremophiles” e.g. thermophiles (thermal vents, hot springs; and halophiles (extreme salt).

UC Berkeley link Archaea Morphology

- By Franciscosp2 - self-made (using xfig), Public Domain, https://commons.wikimedia.org/w/index.php?curid=2652276



Branched isoprene chains

archaea

2. Ether linkages

Mirror image glycerols

Unbranched fatty acids

bacteria

6. Ester linkages

bacteria

archaea

Polymer of sugar and amino acids

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) BACTERIAL CELL

(b) ARCHAEAL CELL

(c) Ester-bonded phospholipid

(d) Ether-bonded phospholipid

a: © Linda Graham: b: © Eye of Science/Photo Researchers, Inc.

Extremophiles

- Can occupy habitats with **extreme conditions**
    - ❑ High salt content – halophiles
    - ❑ Acidity
    - ❑ High methane levels
    - ❑ High temperatures – hyperthermophiles
    - ❑ High radiation

- ex: *Methanopyrus*
    - ❑ Grows in **deep-sea thermal vents** at 98°C (208°F)
    - ❑ Cannot grow at temps under 84°C

- ex: *Sulfolobus* grows in **acidic hot springs** at pH 3

Extremophiles and Their Preferred Conditions

Extremophile Type	Conditions for Optimal Growth
Acidophiles	pH 3 or below
Alkaliphiles	pH 9 or above
Thermophiles	Temperature 60–80°C (140–176°F)
Hyperthermophiles	Temperature 80–122°C (176–250°F)
Psychrophiles	Temperature of -15 -10°C (5-14°F) or lower
Halophiles	Salt concentration of at least 0.2 M
Osmophiles	High sugar concentration

Figure 22.5



- *Deinococcus radiodurans*, visualized in this false color transmission electron micrograph, is a prokaryote that can tolerate very high doses of ionizing radiation. It has developed DNA repair mechanisms that allow it to reconstruct its chromosome even if it has been broken into hundreds of pieces by radiation or heat. (credit: modification of work by Michael Daly; scale-bar data from Matt Russell)

“Heat-loving” (Thermophilic) prokaryotes

Extreme halophiles

Figure 22.6



(a) The Dead Sea is hypersaline (10X salt of seawater, 40x magnesium). Nevertheless, salt-tolerant archaea-bacteria thrive in this sea.

(b) These halobacteria cells can form salt-tolerant bacterial mats.

(credit a: Julien Menichini; credit b: NASA; scale-bar data from Matt Russell)

Some differences between Bacteria and Archaea

1. Bacteria cell walls have peptidoglycan, archaeal cell walls don’t.

2. Bacteria have one ribosomal RNA polymerase (rRNA), archaea and eukaryotes have 3.

3. Bacterial can’t live above 100^{o}C, archaea are extremophiles.

4. Bacteria growth inhibited by antibiotics, archaea unaffected by antibiotics.

Domain Bacteria

- **~50 bacterial phyla (perhaps)**
    - ❑Structural and metabolic features of half or more are still unknown
- Most bacteria favor **moderate conditions**
- Many form **symbiotic** relationships with eukaryotes

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Your microbial fauna:

100 to 200 species in your mouth

128 species in your lungs

up to 1000 species on your skin

300 to 1000 species in your guts



Your microbial fauna:

100 to 200 species in your mouth

128 species in your lungs





The New York Times

July 13, 2010

Microbiomes

A project to gather and sequence DNA from the thousands of microbe species living in and on the human body has completed work on 369 species to date.

Line tips represent bacterial species from genetic surveys. Lines show relationships among species. Only a few had been sequenced.

Circles highlight the species that have been sequenced by the project.

Source: Human Microbiome Project

THE NEW YORK TIMES

Cyanobacteria

- **Photosynthetic bacteria** abundant in fresh water, oceans and wetlands and on surfaces of arid soils
- Named for **blue-green** or cyan color
- The only prokaryotes that **generate oxygen** as a product of photosynthesis
- Gave rise to **plastids** of eukaryotic algae and plants

Cyanobacteria/chloroplast connection



By Kelvinsong - Own work, CC BY-SA 3.0,
https://commons.wikimedia.org/w/index.php?curid=24264654

Cyanobacteria

- Display the greatest **structural diversity** found among bacterial phyla
  - ❑Single cells (unicells) or colonies
  - ❑Filaments
- **Essential ecological roles** in producing organic carbon, O_2 and fixing nitrogen
  - ❑Several kinds form nuisance growths – blooms or biofilms

Cyanobacteria: *Gloeothece* (top left), *Nostoc* (top right), *Calothrix* (bottom left), *Fischerella* (bottom right)



A bloom of cyanobacteria



Biofilms

- Prokaryotes live in communities
- Biofilm is microbial community held together in polysaccharide matrix secreted by the bacteria. These attach to surfaces.
  - Clog pipes
  - Bacterial contamination in food
  - Kitchen counters, cutting boards, sinks, toilets
  - On your teeth

Biofilm development Fig. 22.8



- Five stages of biofilm development are shown. During stage 1, initial attachment, bacteria adhere to a solid surface via weak van der Waals interactions. During stage 2, irreversible attachment, hairlike appendages called pili permanently anchor the bacteria to the surface. During stage 3, maturation I, the biofilm grows through cell division and recruitment of other bacteria. An extracellular matrix composed primarily of polysaccharides holds the biofilm together. During stage 4, maturation II, the biofilm continues to grow and takes on a more complex shape. During stage 5, dispersal, the biofilm matrix is partly broken down, allowing some bacteria to escape and colonize another surface. Micrographs of a *Pseudomonas aeruginosa* biofilm in each of the stages of development are shown. (credit: D. Davis, Don Monroe, PLoS)

Horizontal gene transfer

- Also known as lateral gene transfer
- Movement of one or more genes from one species to another
  - ❑Contrasts with vertical gene transfer from parent to progeny
- Increases genetic diversity
- Common among archaea and bacteria
- Can result in large genetic changes



1 A prokaryote grows in size and develops infoldings in its cell membrane to increase its **surface area to volume ratio.**

2 The infoldings eventually pinch off from the cell membrane, forming an early **endomembrane system.** It encloses the nucleoid, making a **membrane-bound nucleus.** This is the first **eukaryote.**

3 An **aerobic** (oxygen using) **proteobacterium** enters the eukaryote, either as prey or a parasite, and manages to avoid digestion. It becomes an **endosymbiont,** or a cell living inside another cell.

4 The aerobe's ability to use oxygen to make energy becomes an asset for the host, allowing it to thrive in an increasingly oxygen-rich environment as the other eukaryotes go extinct. The proteobacterium is eventually assimilated and becomes a **mitochondrion.**

5 Some eukaryotes go on to acquire additional endosymbionts—the **cyanobacteria,** a group of bacteria capable of photosynthesis. They become **chloroplasts.**

Horizontal gene transfer

At least 17% of the genes present in the common human gut inhabitant *E. coli* came from other bacteria

About 80% of prokaryotic genes have been involved in horizontal transfer at some point in their history

Important concepts

- Bacteria and Archaea evolved from a common ancestor
- Eukaryotic nucleus and cytoplasm likely arose in an ancient archaeal organism
- Mitochondria and plastids originated from 2 phlya: proteobacteria and cyanobacteria by endosymbiosis
  - ❑Transfer of many genes from bacteria to eukaryotes

Proteobacteria

AMAZING DIVERSITY OF FORM AND METABOLISM

FIVE MAJOR SUBGROUPS – CALLED Α, Β, Γ, Δ, Ε

Proteobacteria (all gram negative)

α-proteobacteria (Alpha)	• **Ancestors of mitochondria**, *Rhizobium* (beneficial plant associations), *Agrobacterium* (plant tumors)
β-proteobacteria (*Beta)	• *Nitrosomonas* (nitrogen fixing)
γ-proteobacteria (Gamma)	• *Neisseria* (gonorrhea), *Vibrio* (**cholera**), ***Salmonella***, *Escherichia coli* (***E. coli***)
δ -proteobacteria (Delta)	• Myxobacteria (colonial), bdellovibrios (predatory)
ε –proteobacteria (Epsilon)	• ***Helicobacter*** (stomach ulcers)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Nigel Cattlin/Visuals Unlimited

Agrobacterium tumifasciens infection

Phylum Proteobacteria



Bacteria of Phylum Proteobacteria		
Class	**Representative organisms**	**Representative micrograph**
Alpha Proteobacteria Some species are photoautotrophic but some are symbionts of plants and animals and others are pathogens. Eukaryotic mitochondria are thought be derived from bacteria in this group.	*Rhizobium* Nitrogen-fixing endosymbiont associated with the roots of legumes *Rickettsia* Obligate intracellular parasite that causes typhus and Rocky Mountain Spotted Fever (but not rickets, which is caused by Vitamin C deficiency)	5 µm *Rickettsia rickettsia,* stained red, grow inside a host cell.
Beta Proteobacteria This group of bacteria is diverse. Some species play an important role in the nitrogen cycle.	*Nitrosomas* Species from this group oxidize ammonia into nitrite. *Spirillum minus* Causes rat-bite fever	1 µm *Spirillum minus*
Gamma Proteobacteria Many are beneficial symbionts that populate the human gut, but others are familiar human pathogens. Some species from this subgroup oxidize sulfur compounds.	*Escherichia coli* Normally beneficial microbe of the human gut, but some strains cause disease *Salmonella* Certain strains cause food poisoning or typhoid fever *Yersinia pestis* Causative agent of Bubonic plague *Psuedomonas aeruginosa* Causes lung infections *Vibrio cholera* Causative agent of cholera *Chromatium* Sulfur-producing bacteria that oxidize sulfur, producing H_2S	1 µm *Vibrio cholera*
Delta Proteobacteria Some species generate a spore-forming fruiting body in adverse conditions. Others reduce sulfate and sulfur.	*Myxobacteria* Generate spore-forming fruiting bodies in adverse conditions *Desulfovibrio vulgaris* Aneorobic, sulfate-reducing bacterium	500 nm *Desulfovibrio vulgaris*
Epsilon Proteobacteria Many species inhabit the digestive tract of animals as symbionts or pathogens. Bacteria from this group have been found in deep-sea hydrothermal vents and cold seep habitats.	*Campylobacter* Causes blood poisoning and intestinal inflammation *Heliobacter pylori* Causes stomach ulcers	500 nm *Campylobacter*

Five of the Major Clades of Bacteria



Copyright © 2005 Pearson Education, Inc. Publishing as Pearson Benjamin Cummings. All rights reserved.

Other bacterial phyla

Bacteria: Chlamydia, Spirochaetae, Cyanobacteria, and Gram-positive

Phylum	Representative organisms	Representative micrograph
Chlamydias All members of this group are obligate intracellular parasites of animal cells. Cells walls lack peptidoglycan.	*Chlamydia trachomatis* Common sexually transmitted disease that can lead to blindness	 In this pap smear, *Chlamydia trachomatis* appear as pink inclusions inside cells.
Spirochetes Most members of this species, which has spiral-shaped cells, are free-living aneaerobes, but some are pathogenic. Flagella run lengthwise in the periplasmic space between the inner and outer membrane.	*Treponema pallidum* Causative agent of syphilis *Borrelia burgdorferi* Causative agent of Lyme disease	 *Treponema pallidum*
Cyanobacteria Also known as blue-green algae, these bacteria obtain their energy through photosynthesis. They are ubiquitous, found in terrestrial, marine, and freshwater environments. Eukaryotic chloroplasts are thought be derived from bacteria in this group.	*Prochlorococcus* Believed to be the most abundant photosynthetic organism on earth; responsible for generating half the world's oxygen	 *Phormidium*
Gram-positive Bacteria Soil-dwelling members of this subgroup decompose organic matter. Some species cause disease. They have a thick cell wall and lack an outer membrane.	*Bacillus anthracis* Causes anthrax *Clostridium botulinum* Causes Botulism *Clostridium difficile* Causes diarrhea during antibiotic therapy *Streptomyces* Many antibiotics, including streptomyocin, are derived from these bacteria. *Mycoplasmas* These tiny bacteria, the smallest known, lack a cell wall. Some are free-living, and some are pathogenic.	 *Clostridium difficile*

Structure and Movement

- Bacteria and archaea share
  - small size
  - rapid growth
  - simple cellular structure
- Bacteria and archaea are just **1–5 µm** in diameter
  - ❑Plant and animal cells - between 10 and 100 µm in diameter
- Small cell size less cellular material BUT **faster cell division**

Prokaryotes and a eukaryotic cell



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Prokaryotic cell



Cell shape and arrangement

- Five major shapes
  - ❑ Spheres – **cocci**
  - ❑ Rods – **bacilli**
  - ❑ Comma-shaped – **vibrios**
  - ❑ Spiral-shaped – **spirochaetes** are flexible, while **spirilli** are rigid
- Occur as single cells, pairs, or filaments

The most common shapes of bacteria (colorized SEM)


2 µm
Cocci
(coccus)
Vibrio cholerae
Copyright © 2001 Dennis Kunkel Microscopy, Inc. / Dennis Kunkel





Bacteria cell-wall structure

- Most have rigid cell wall outside the plasma membrane
  - Maintain **cell shape** and help **protect against attack**
  - Also help avoid lysis in hypotonic solutions
- Archaea and some bacteria use protein
- Most bacteria use **peptidoglycan** (polysaccharide chains cross-linked with unusual peptides)

Gram stain

- **Gram positive**
  - ❑ Relatively **thick** peptidoglycan layer
  - ❑ Purple dye held in thick layer
  - ❑ Cells are stained **purple**
  - ❑ **Vulnerable to penicillin** that interferes in cell wall synthesis

- **Gram negative**
  - ❑ Less peptidoglycan and a thin outer envelope of lipopolysaccharides
  - ❑ Lose purple stain but retain final pink stain
  - ❑ Cells are stained **pink**
  - ❑ **Resists penicillin** and requires other antibiotics

Gram-positive and gram-negative bacteria



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Gram-positive and gram-negative bacteria



(a) Gram-positive



(b) Gram-negative

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Acidic polysaccharides

Thick peptidoglycan layer

Plasma membrane

(a) Gram-positive: thick peptidoglycan layer, no outer envelope



Lipopolysaccharide-rich outer envelope

Thin peptidoglycan layer

Plasma membrane

(b) Gram-negative: thinner peptidoglycan layer, with outer envelope

- All gram positive bacteria are in one phyla.
- Gram negative bacteria are in many phyla, including Proteobacteria, Chlamydias, Cyanobacteria, Spirochetes, and others.

Motility (ability to move)

- Allows cells to move to favorable conditions
- Respond to chemical signals
- Swim, twitch, glide or adjust flotation

Flagella

- Swimming
- Different from eukaryotic flagella
- Like an outboard motor
- Differ in number and location of flagella

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



The motor rotates the hook, which links the internal motor to the external filament, composed of a protein known as flagellin.

A set of protein rings anchors the flagellum in the plasma membrane and cell wall. As protons (H^+) flow into the cell through membrane proteins within the motor, these proteins change conformation, thereby turning the rings.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) Bacteria with a single short flagellum



(b) Bacterium with multiple long flagella

a: © Dennis Kunkel Microscopy/Phototake; b: © Scientifica/Visuals Unlimited

Pili (pilus)

- Twitch or glide across surfaces
- Attach to surfaces (biofilms)
- Threadlike structures on surface of cell
- Also play important roles in bacterial reproduction and disease processes





Prokaryotic reproduction and new genetic combinations

- Binary fission – continual synthesis of DNA
- Transformation – take up of genes from the environment
- Conjugation – direct transfer of genes between prokaryotes
- Transduction – transfer of genes by viruses



Prokaryotic conjugation



- **Binary fission** – divide by splitting in two
- Basis for widely used method of detecting and counting bacteria in samples
  - ❑Place measured volume of sample into plastic dishes of agar
  - ❑Each single cell will form a visible colony
- Can also use fluorescent dye that binds bacterial DNA to directly count bacteria



(a) Bacterium undergoing binary fission



(b) Colonies developed from single cells

Nutrition and Metabolism

- More diverse types of metabolism than any other group of organisms
- Can be classified by
  - ☐ Nutrition
  - ☐ Response to oxygen
  - ☐ Presence of specialized metabolic processes



Table 27.1 Major Nutritional Modes

Mode of Nutrition	Energy Source	Carbon Source	Types of Organisms
Autotroph			
Photoautotroph	Light	CO_2	Photosynthetic prokaryotes, including cyanobacteria; plants; certain protists (algae)
Chemoautotroph	Inorganic chemicals	CO_2	Certain prokaryotes (for example, *Sulfolobus*)
Heterotroph			
Photoheterotroph	Light	Organic compounds	Certain prokaryotes
Chemoheterotroph	Organic compounds	Organic compounds	Many prokaryotes and protists; fungi; animals; some parasitic plants

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Classification by oxygen response

- **Obligate aerobes** – require oxygen
- **Facultative anaerobes** – can use oxygen or not
- **Obligate anaerobes** – cannot tolerate oxygen

Classification by special metabolism

- **Diazotrophs** – conduct nitrogen fixation
  - Enzyme nitrogenase converts inorganic nitrogen gas (N_2) into ammonia (NH_3)
  - Plants and eukaryotic algae depend on ammonia to make nitrogen containing compounds
  - *Rhizobium* (proteobacterium)
  - Heterocysts – specialized cells for nitrogen fixation in cyanobacteria like *Anabaena*



Ecological Roles and Biotechnological Applications

- Carbon, nitrogen, phosphorous cycles
  - ❑Producers synthesize organic compounds used by other organisms as food
  - ❑Decomposers (saprobes) break down dead organisms to release minerals for reuse

THE CARBON CYCLE



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

THE NITROGEN CYCLE


Nitrogen in atmosphere (N_2)
Plants
Assimilation
Denitrifying bacteria
Nitrogen-fixing bacteria in root nodules of legumes
Decomposers (aerobic and anaerobic bacteria and fungi)
Nitrates (NO_3^-)
Nitrifying bacteria
Ammonification
Nitrification
Ammonium (NH_4^+)
Nitrites (NO_2^-)
Nitrogen-fixing soil bacteria
Nitrifying bacteria

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

THE PHOSPHOROUS CYCLE



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Symbiotic roles

- **Symbiosis**
  - An organism that lives in close association with one or more other organisms
- **Parasitism**
  - One partner benefits at the expense of the other
- **Mutualism**
  - Association is beneficial to both partners

Abyssopelagic zone



Mutualism: bacterial “headlights”
Lanterneye or flashlight fish



Anglerfish female (with male)


Symbiotic bacteria make light

- **Human microbiome**
  - On human skin and in our digestive and reproductive systems
  - Provide services using traits that humans do not possess
  - Humans and their microbiome are **superorganisms**
    - Diverse types of bacterial metabolism have co-evolved with humans
    - The human microbiome
    - microbiome

- **Pathogens**
  - Organisms that obtain organic compounds from living hosts and cause disease (abnormal functioning of host)
  - Pathogen – parasitic microbe **causes disease** symptoms
  - Examples of prokaryote caused disease:
    - Cholera, leprosy, tetanus, pneumonia, whooping cough, diphtheria, Lyme disease, scarlet fever, rheumatic fever, typhoid fever, bacterial dysentery, and tooth decay (very incomplete list)

Lyme disease, a bacterial disease transmitted by ticks



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.



Putting prokaryotes to work in sewage treatment facilities



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Bioremediation for an oil spill